

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



4-Hydroxyphenylpropionylglycine

Item No. 36389

CAS Registry No.: 3850-43-9

Formal Name: N-[3-(4-hydroxyphenyl)-1-oxopropyl]-glycine

Synonyms: Phloretic Acid Glycine Conjugate,

Phloretylglycine, 4-hydroxy PPG

MF: $C_{11}H_{13}NO_4$ 223.2 FW: ≥95% **Purity:**

UV/Vis.: λ_{max} : 226 nm

Supplied as: A solid Storage: -20°C Stability: ≥4 years Item Origin: Synthetic

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

4-Hydroxyphenylpropionylglycine is supplied as a solid. A stock solution may be made by dissolving the 4-hydroxyphenylpropionylglycine in the solvent of choice, which should be purged with an inert gas. 4-Hydroxyphenylpropionylglycine is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF). The solubility of 4-hydroxyphenylpropionylglycine in ethanol is approximately 20 mg/ml and approximately 10 mg/ml in DMSO and DMF.

Description

4-Hydroxyphenylpropionylglycine is a metabolite of the conditionally essential amino acid tyrosine.¹ It is formed from tyrosine via aromatic amino acid aminotransferase, tyrosine aminotransferase, and gut microbiota, followed by glycine conjugation. 4-Hydroxyphenylpropionylglycine is also a metabolite of the phenol phloretin (Item No. 14452).²

References

- 1. Anesi, A., Berding, K., Clarke, G., et al. Metabolomic workflow for the accurate and high-throughput exploration of the pathways of tryptophan, tyrosine, phenylalanine, and branched-chain amino acids in human biofluids. J. Proteome Res. 21(5), 1262-1275 (2022).
- 2. Monge, P., Solheim, E., and Scheline, R.R. Dihydrochalcone metabolism in the rat: Phloretin. Xenobiotica 14(2), 917-924 (1984).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 11/02/2023

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM