



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)


# PRODUCT INFORMATION



## 2-Methylglutaric Acid

Item No. 36440

**CAS Registry No.:** 617-62-9  
**Formal Name:** 2-methyl-pentanedioic acid  
**Synonyms:**  $\alpha$ -Methylglutaric Acid,  
NSC 5941, NSC 408456  
**MF:** C<sub>6</sub>H<sub>10</sub>O<sub>4</sub>  
**FW:** 146.1  
**Purity:**  $\geq$ 95%  
**Supplied as:** A solid  
**Storage:** -20°C  
**Stability:**  $\geq$ 4 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

### Laboratory Procedures

2-Methylglutaric acid is supplied as a solid. A stock solution may be made by dissolving the 2-methylglutaric acid in the solvent of choice, which should be purged with an inert gas. 2-Methylglutaric acid is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF). The solubility of 2-methylglutaric acid in DMSO is approximately 16 mg/ml and approximately 20 mg/ml in DMF. It is slightly soluble in ethanol.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of 2-methylglutaric acid can be prepared by directly dissolving the solid in aqueous buffers. The solubility of 2-methylglutaric acid in PBS (pH 7.2) is approximately 10 mg/ml. We do not recommend storing the aqueous solution for more than one day.

### Description

2-Methylglutaric acid is a dicarboxylic acid and derivative of the L-leucine metabolite 3-methylglutaric acid (Item No. 34357) that has been found in *K. excelsa* honey.<sup>1</sup> 2-Methylglutaric acid levels are decreased by the methylation inhibitor 5-aza-2'-deoxycytidine (decitabine; Item No. 11166) in several bladder cancer cell lines.<sup>2</sup>

### References

1. Wilkins, A.L., Lu, Y., and Tan, S.-T. Extractives from New Zealand honeys. 5. Aliphatic dicarboxylic acids in New Zealand rewarewa (*Knightea excelsa*) honey. *J. Agric. Food Chem.* **43**, 3021-3025 (1995).
2. Ambati, C.R., Vantaku, V., Donepudi, S.R., et al. Measurement of methylated metabolites using Liquid Chromatography-Mass Spectrometry and its biological application. *Anal. Methods* **11(1)**, 49-57 (2019).

#### WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 03/02/2023

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA

**PHONE:** [800] 364-9897  
[734] 971-3335

**FAX:** [734] 971-3640

CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM