

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



## Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



# PRODUCT INFORMATION



### Steviolbioside

Item No. 36513

CAS Registry No.: 41093-60-1

(4α)-13-[(2-O-β-D-glucopyranosyl-β-D-glucopyranoFormal Name:

glucopyranosyl)oxy]-kaur-16-en-18-oic acid


Synonym: Steviobioside

MF:  $C_{32}H_{50}O_{13}$ 642.7 FW: ≥98% **Purity:** Supplied as: A solid

Storage: -20°C Stability: ≥4 years

Item Origin: Plant/Stevia rebaudiana

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



#### **Laboratory Procedures**

Steviolbioside is supplied as a solid. A stock solution may be made by dissolving the steviolbioside in the solvent of choice, which should be purged with an inert gas. Steviolbioside is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF). The solubility of steviolbioside in ethanol is approximately 5 mg/ml and approximately 25 mg/ml in DMSO and DMF.

#### Description

Steviolbioside is a natural non-caloric sweetener and glycoside that has been found in S. rebaudiana leaves and an active metabolite of rebaudioside A (Item No. 11894). It inhibits the proliferation of MDA-MB-231 breast cancer cells when used at a concentration of 250 µg/ml. Steviolbioside is active against M. tuberculosis (MIC =  $3.75 \mu g/ml$ ).<sup>2</sup>

#### References

- 1. Chen, J.-M., Ding, L., Sui, X.-C., et al. Production of a bioactive sweetener steviolbioside via specific hydrolyzing ester linkage of stevioside with a β-galactosidase. Food Chem. 196, 155-160 (2016).
- 2. Sharipova, R.R., Strobykina, I.Y., Mordovskoi, G.G., et al. Antituberculosis activity of glycosides from Stevia rebaudiana and hybrid compounds of steviolbioside and pyridinecarboxylic acid hydrazides. Chem. Nat. Compd. 46, 902-905 (2011).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

## WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 06/01/2023

#### **CAYMAN CHEMICAL**

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM