

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



## Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



# PRODUCT INFORMATION



## **Pyrimorph**

Item No. 37286

CAS Registry No.: 868390-90-3

Formal Name: 3-(2-chloro-4-pyridinyl)-3-[4-

(1,1-dimethylethyl)phenyl]-1-(4-

morpholinyl)-2-propen-1-one


MF: C22H25CIN2O2

384.9 FW: ≥90% **Purity:** 

UV/Vis.:  $\lambda_{max}$ : 244 nm Supplied as: A solid

Storage: -20°C Stability: ≥4 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



## **Laboratory Procedures**

Pyrimorph is supplied as a solid. A stock solution may be made by dissolving the pyrimorph in the solvent of choice, which should be purged with an inert gas. Pyrimorph is soluble in the organic solvent dimethyl formamide at a concentration of approximately 1 mg/ml.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of pyrimorph can be prepared by directly dissolving the solid in aqueous buffers. The solubility of pyrimorph in PBS (pH 7.2) is approximately 0.20 mg/ml. We do not recommend storing the aqueous solution for more than one day.

#### Description

Pyrimorph is a fungicide.<sup>1</sup> It is active against the plant pathogenic fungi P. infestans, P. capsica, and R. solani (EC<sub>50</sub>s = 1.36, 0.72, and 4.44  $\mu$ g/ml, respectively). Pyrimorph is toxic to zebrafish (LC<sub>50</sub> = 19.8 mg/L).2

#### References

- 1. Chen, X., Yuan, H., Qin, Z., et al. Preliminary studies on antifungal activity of pyrimorph. Nongyaoxue Xuebao 9(3), 229-234 (2007).
- 2. Zhao, C., Liu, B., Wang, J., et al. Acute toxicity and bioconcentration of pyrimorph in zebrafish, Brachydanio rerio. Pest. Manag. Sci. 67(9), 1178-1183 (2011).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 07/12/2023

## **CAYMAN CHEMICAL**

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM