

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



CFMTI

Item No. 37546

CAS Registry No.: 864864-17-5

Formal Name: 2-cyclopropyl-5-[1-(2-fluoro-3-

≥4 years

pyridinyl)-5-methyl-1H-1,2,3-triazol-4-

yl]-2,3-dihydro-1H-isoindol-1-one

MF: C₁₉H₁₆FN₅O

349.4 FW: ≥90% **Purity:**

 λ_{max} : 272 nm UV/Vis.: Supplied as: A solid Storage: -20°C

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

CFMTI is supplied as a solid. A stock solution may be made by dissolving the CFMTI in the solvent of choice, which should be purged with an inert gas. CFMTI is soluble in organic solvents such as chloroform, DMSO, and dimethyl formamide (DMF). The solubility of CFMTI in chloroform and DMF is approximately 1 and 10 mg/ml, respectively. CFMTI is slightly soluble in DMSO.

Description

Stability:

CFMTI is a metabotropic glutamate receptor 1 (mGluR1) allosteric antagonist (IC $_{50}$ = 2.6 nM in CHO cells expressing the human receptor). 1.2 It is selective for mGluR1 over mGluR5 by greater than 2,000-fold, as well as over mGluR2, -4, -6, -7, or -8 at 10 μM in CHO cells expressing the human receptors.² CFTMI reduces methamphetamine-induced hyperlocomotion in mice when administered orally at doses ranging from 1 to 10 mg/kg.

References

- 1. Ito, S., Hirata, Y., Nagatomi, Y., et al. Discovery and biological profile of isoindolinone derivatives as novel metabotropic glutamate receptor 1 antagonists: A potential treatment for psychotic disorders. Bioorg. Med. Chem. Lett. 19(18), 5310-5313 (2009).
- 2. Satow, A., Suzuki, G., Maehara, S., et al. Unique antipsychotic activities of the selective metabotropic glutamate receptor 1 allosteric antagonist 2-cyclopropyl-5-[1-(2-fluoro-3-pyridinyl)-5-methyl-1H-1,2,3triazol-4-yl]-2,3-dihydro-1H-isoindol-1-one. J. Pharmacol. Exp. Ther. 330(1), 179-190 (2009).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information Buyer agrees to purchase the mater can be found on our website.

Copyright Cayman Chemical Company, 02/27/2023

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM