

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



## Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



# PRODUCT INFORMATION



### Histone H4 (human, recombinant; His-tagged)

Item No. 38257

#### **Overview and Properties**

Source: Recombinant human N-terminal His-tagged histone H4 expressed in E. coli

**Amino Acids: Uniprot No.:** P62805 Molecular Weight: 13.4 kDa

-80°C (as supplied) Storage:

Stability: ≥6 months

batch specific (≥80% estimated by SDS-PAGE) **Purity:** 


Supplied in: A solution in water

**Protein** 

batch specific mg/ml Concentration:

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

#### **Image**



Lane 1: MW Markers Lane 2: Histone H4 (2 µg) Lane 2: Histone H4 (4 µg)

SDS-PAGE Analysis of Histone H4.

Representative gel image shown; actual purity may vary between each batch.

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 05/22/2023

**CAYMAN CHEMICAL** 

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM

## **PRODUCT INFORMATION**



#### Description

Histone H4 is one of four core histone proteins that are involved in the organization of DNA into chromatin. Histones are globular proteins with unstructured N-terminal tails and are subject to a variety of post-translational modifications (PTMs), such as methylation, acetylation, phosphorylation, and citrullination, that can influence chromatin structure and regulate gene transcription. Histone H4 PTMs function as epigenetic regulators of transcription by affecting chromatin structure and enhancing transcription factor binding, as well as play a key role in the maintenance of genomic integrity. Levels of histone H4 are increased in the inferior and middle temporal gyrus in postmortem brains from patients with Alzheimer's disease.

#### References

- 1. Wang, Y., Li, M., Stadler, S., et al. Histone hypercitrullination mediates chromatin decondensation and neutrophil extracellular trap formation. J. Cell Biol. 184(2), 205-213 (2009).
- 2. Hyun, K., Jeon, J., Park, K., et al. Writing, erasing and reading histone lysine methylations. Exp. Mol. Med. 49(4), e324 (2017).
- Vettese-Dadey, M., Grant, P.A., Hebbes, T.R., et al. Acetylation of histone H4 plays a primary role in enhancing transcription factor binding to nucleosomal DNA in vitro. EMBO J. 15(10), 2508-2518 (1996).
- 4. Jørgensen, S., Schotta, G., and Sørensen, C.S. Histone H4 Lysine 20 methylation: Key player in epigenetic regulation of genomic integrity. *Nucleic Acids Res.* **41(5)**, 2797-2806 (2013).
- 5. Narayan, P.J., Lill, C., Faull, R., et al. Increased acetyl and total histone levels in post-mortem Alzheimer's disease brain. *Neurobiol. Dis.* **74**, 281-294 (2015).