



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



PRODUCT INFORMATION

1,2,3,4,7,8-Hexachlorodibenzofuran

Item No. 38619

CAS Registry No.: 70648-26-9

Synonyms: 1,2,3,4,7,8-HxCDF, PCDF 118

MF: C₁₂H₂Cl₆O


FW: 374.9

Purity: ≥90%

Supplied as: A solid

Storage: -20°C

Stability: ≥4 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

1,2,3,4,7,8-Hexachlorodibenzofuran is supplied as a solid. A stock solution may be made by dissolving the 1,2,3,4,7,8-hexachlorodibenzofuran in the solvent of choice, which should be purged with an inert gas. 1,2,3,4,7,8-Hexachlorodibenzofuran is slightly soluble in chloroform (warmed).

Description

1,2,3,4,7,8-Hexachlorodibenzofuran is a dioxin-like polychlorinated dibenzofuran (PCDF).¹ It induces the expression of the genes encoding the cytochrome P450 (CYP) isoforms CYP1A1 and CYP1B1, as well as the aryl hydrocarbon receptor repressor (AhRR) in primary human peripheral blood lymphocytes and increases ethoxresorufin-O-deethylase (EROD) activity, a marker of CYP1A1 activity, in isolated human peripheral blood lymphocytes in a concentration-dependent manner. 1,2,3,4,7,8-Hexachlorodibenzofuran (300, 600, and 1,000 µg/kg) increases the percentage of mouse fetuses with cleft palates or hydronephrosis in pregnant dams.² It has been found in human breastmilk, freshwater fish, and in the air near municipal waste incinerators.³⁻⁵

References

- van Ede, K.I., Gaisch, K.P.J., van den Berg, M., et al. Differential relative effect potencies of some dioxin-like compounds in human peripheral blood lymphocytes and murine splenic cells. *Toxicol. Lett.* **226**(1), 43-52 (2014).
- Birnbaum, L.S., Harris, M.W., Barnhart, E.R., et al. Teratogenicity of three polychlorinated dibenzofurans in C57BL/6N mice. *Toxicol. Appl. Pharmacol.* **90**(2), 206-216 (1987).
- Takekuma, M., Saito, K., Ogawa, M., et al. Levels of PCDDs, PCDFs and Co-PCBs in human milk in Saitama, Japan, and epidemiological research. *Chemosphere* **54**(1), 127-135 (2004).
- Tawara, K., Nishijo, M., Honda, R., et al. Effects of maternal dioxin exposure on newborn size at birth among Japanese mother-infant pairs. *Environ. Health Prev. Med.* **14**(2), 88-95 (2009).
- Hu, S.-W., ChangChien, G.-P., and Chan, C.-C. PCDD/Fs levels in indoor environments and blood of workers of three municipal waste incinerators in Taiwan. *Chemosphere* **55**(4), 611-620 (2004).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 04/27/2023

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 - USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM