



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION



Soluble Epoxide Hydrolase PROTAC 1a

Item No. 38876

Formal Name: N-(1-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxisoindolin-4-yl)oxy)-2-oxo-6,9,12,15-tetraoxa-3-azaheptadecan-17-yl)-4-(((1r,4r)-4-(3-(4-(trifluoromethoxy)phenyl)ureido)cyclohexyl)oxy)benzamide

Synonyms: sEH PROTAC 1a, sEH Proteolysis-targeting Chimera 1a, Soluble Epoxide Hydrolase Proteolysis-targeting Chimera 1a

MF: C₄₆H₅₃F₃N₆O₁₄

FW: 971.0


Purity: ≥95%

UV/Vis.: λ_{max}: 245 nm

Supplied as: A solid

Storage: -20°C

Stability: ≥3 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Description

Soluble epoxide hydrolase (sEH) PROTAC 1a is a proteolysis-targeting chimera (PROTAC) that contains cereblon ligand 1 (Item No. 21589) connected to the sEH inhibitor *t*-TUCB (Item No. 36384) by a linker region.¹ It drives sEH degradation and selectively inhibits the hydrolase activity of sEH (IC₅₀ = 0.8 nM) over its phosphatase activity (IC₅₀ = >10,000 nM). sEH PROTAC 1a is also selective for the degradation of cytosolic over peroxisomal sEH and drives its degradation *via* the lysosome rather than the proteasome. It reduces thapsigargin-induced increases in phosphorylated inositol-requiring enzyme 1α (IRE1α) levels and X-box binding protein 1 (XBP1) splicing in HepG2 and 293T cells, indicating a reduction in ER stress.

Reference

1. Wang, Y., Morisseau, C., Takamura, A., *et al.* PROTAC-Mediated selective degradation of cytosolic soluble epoxide hydrolase enhances ER stress reduction. *ACS Chem. Biol.* **18**(4), 884-896 (2023).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 07/12/2023

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM

WWW.CAYMANCHEM.COM