



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



PRODUCT INFORMATION



Cnicin

Item No. 38969

CAS Registry No.: 24394-09-0

Formal Name: (3R)-3,4-dihydroxy-2-methylenebutanoic acid, (3aR,4S,6E,10Z,11aR)-2,3,3a,4,5,8,9,11a-octahydro-10-(hydroxymethyl)-6-methyl-3-methylene-2-oxocyclodeca[b]furan-4-yl ester

Synonym: (+)-Cnicin

MF: C₂₀H₂₆O₇

FW: 378.4


Purity: ≥95%

Supplied as: A solid

Storage: -20°C

Stability: ≥4 years

Item Origin: Plant/Centaurea maculosa



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Cnicin is supplied as a solid. A stock solution may be made by dissolving the cnicin in the solvent of choice, which should be purged with an inert gas. Cnicin is soluble in organic solvents such as acetonitrile and methanol.

Description

Cnicin is a germacranolide sesquiterpene lactone that has been found in *Centaurea* and has diverse biological activities.¹⁻³ It irreversibly inhibits the UDP-N-acetylglucosamine (UDP-GlcNAc) 1-carboxyvinyltransferase MurA (IC_{50} s = 16.7 and 10.3 for the *E. coli* and *P. aeruginosa* enzymes, respectively) and is active against *B. cereus* and *M. flavus* (MICs = 1.25 and 5 µg/ml, respectively).^{1,2} Cnicin inhibits the growth of MCF-7 breast and OVCAR-3 ovarian cancer cells (GI_{50} s = 37.8 and 37.4 µM, respectively).² It reduces snake venom-induced paw edema in rats when administered at doses ranging from 2.5 to 10 mg/kg.³

References

1. Bachelier, A., Mayer, R., and Klein, C.D. Sesquiterpene lactones are potent and irreversible inhibitors of the antibacterial target enzyme MurA. *Bioorg. Med. Chem. Lett.* **16**(21), 5605-5609 (2006)
2. Saroglou, V., Karioti, A., Demetzos, C., et al. Sesquiterpene lactones from *Centaurea spinosa* and their antibacterial and cytotoxic activities. *J. Nat. Prod.* **68**(9), 1404-1407 (2005).
3. Demiroz, T., Albayrak, G., Nalbantsoy, A., et al. Anti-inflammatory properties of *Centaurea calolepis* Boiss. and cnicin against *Macrovipera lebetina obtusa* (Dwigubsky, 1832) and *Montivipera xanthina* (Gray, 1849) venoms in rat. *Toxicon* **152**, 37-42 (2018).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 - USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM