

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



COOH

20-HETE-d₆ Item No. 390030

CAS Registry No.: 2548939-89-3

Formal Name: 20-hydroxy-5Z,8Z,11Z,14Z-eicosatetraenoic-

16,16,17,17,18,18-d₄ acid

Synonyms: 20-Hydroxyeicosatetraenoic Acid-d₄,

20-hydroxy Arachidonic Acid-d

MF: $C_{20}H_{26}D_6O_3$ FW:

326.5 **Chemical Purity:** ≥95% (20-HETE)

Deuterium

Incorporation: \geq 99% deuterated forms (d₁-d₆); \leq 1% d₀

A solution in acetonitrile Supplied as:

-20°C Storage: Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

20-HETE-d₆ is intended for use as an internal standard for the quantification of 20-HETE (Item No. 90030) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

20-HETE-d₆ is supplied as a solution in acetonitrile. To change the solvent, simply evaporate the acetonitrile under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as ethanol, DMSO, and dimethyl formamide purged with an inert gas can be used. The solubility of 20-HETE-d₆ in these solvents is approximately 10 mg/ml.

Description

20-HETE is a cytochrome P450 (CYP450) metabolite postulated to play an autacoid role in the renal and cerebral vasculature. In rat cerebral microvessels, 20-HETE is a vasoconstrictor that mediates pressure-induced autoregulatory vasoconstriction.² 20-HETE is excreted mainly as the glucuronide conjugate. The concentration of free 20-HETE (20-40 pg/ml in human urine) is about 10-fold lower than the corresponding concentration of the 20-glucuronide.³

References

- 1. McGiff, J.C. and Quilley, J. 20-HETE and the kidney: Resolution of old problems and new beginnings. Am. J. Physiol. 277(3 Pt 2), R607-R623 (1999).
- Gebremedhin, D., Lange, A.R., Lowry, T.F., et al. Production of 20-HETE and its role in autoregulation of cerebral blood flow. Circ. Res. 87(1), 60-65 (2000).
- Prakash, C., Zhang, J.Y., Falck, J.R., et al. 20-Hydroxyeicosatetraenoic acid is excreted as a glucuronide conjugate in human urine. Biochem. Biophys. Res. Commun. 185(2), 728-733 (1992).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 05/06/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM