

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



C16 dihydro Ceramide-d_o (d18:0/16:0-d_o)

Item No. 39389

CAS Registry No.: 2260669-52-9

N-[(1S,2R)-2-hydroxy-1-(hydroxymethyl)heptadecyl]-Formal Name:

hexadecanamide-13,13,14,14,15,15,16,16,16-d_o

Synonyms: Cer(d18:0/16:0-d_o), Ceramide (d18:0/16:0-d_o),

N-hexadecanoyl-D-erythro-Dihydrosphingosine-do.

N-Palmitoyl Sphinganine-do

MF: $C_{34}H_{60}D_{9}NO_{3}$

FW: 549.0


Chemical Purity: ≥98% (C16 dihydro Ceramide (d18:0/16:0))

Deuterium

 \geq 99% deuterated forms (d₁-d₉); \leq 1% d₀ Incorporation:

Supplied as: A solid -20°C Storage: ≥4 years Stability:

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

C16 dihydro Ceramide-d_o (d18:0/16:0-d_o) is intended for use as an internal standard for the quantification of C16 dihydro ceramide (Item No. 24369) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

C16 dihydro Ceramide-d_o (d18:0/16:0-d_o) is supplied as a solid. A stock solution may be made by dissolving the C16 dihydro ceramide-do (d18:0/16:0-do) in the solvent of choice, which should be purged with an inert gas. C16 dihydro Ceramide-d_o (d18:0/16:0-d_o) is soluble in ethanol and DMSO.

Description

C16 dihydro Ceramide is a bioactive sphingolipid and precursor in the de novo synthesis of C16 ceramide (d18:0/16:0) (Item No. 10681) that lacks the 4,5-trans double bond. C16 dihydro Ceramide (46 nM) inhibits C16 ceramide-induced membrane permeabilization, measured as cytochrome c oxidation, in rat liver mitochondria in a concentration-dependent manner. It also inhibits C16 ceramide-induced channel formation in liposomes. C16 dihydro ceramide is biologically inactive as a single agent, lacking the ability to induce apoptosis, cytochrome c release, or channel formation in phospholipid membranes in the absence of C16 ceramide.

Reference

1. Stiban, J., Fistere, D., and Colombini, M. Dihydroceramide hinders ceramide channel formation: Implications on apoptosis. Apoptosis 11(5), 773-780 (2006).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 02/04/2025

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM