

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Glypican-3 Rabbit Monoclonal Antibody (APC) (Clone 024)

Item No. 39540

Overview and Properties

This vial contains protein A-affinity purified monoclonal antibody Contents:

Synonyms: GPC3, MXR7, OCI-5

Immunogen: A synthetic peptide corresponding to the central region of human glypican-3

Cross Reactivity: (+) Glypican-3 Species Reactivity: (+) Human Form: Liquid

2-8°C (as supplied) Storage:

Stability: ≥1 year


Storage Buffer: PBS with 0.5% BSA and 0.03% ProClin™ 300

Concentration: $10 \mu l/Test$, 0.1 mg/ml

024 Clone: Rabbit Host: Isotype: **IgG**

Application: Flow cytometry (FC); the optimal working concentration/dilution should be determined

Image



Flow cytometric analysis of Human CPC3 expression in HepG2 cells. Cells were stained with Glypican-3 Rabbit Monoclonal Antibody (APC) (Clone 024). The fluorescence histograms were derived from fated evens with the forward and side light-scatter characteristics on intact Clone.

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 11/08/2023

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM

PRODUCT INFORMATION



Description

Glypican-3 (GPC3) is a membrane-bound heparan sulfate proteoglycan.¹ It is composed of a signal peptide for membrane translocation, a cysteine-rich domain that contains a proteolytic cleavage site for proprotein convertases, a stalk region that contains heparan sulfate attachment sites, and a signal sequence for glycosylphosphatidylinositol (GPI) attachment, which anchors it to the cell surface. GPC3 is ubiquitously expressed during embryonic development and is only expressed in select adult tissues, including gastric glands and kidney tubules.^{2,3} It is involved in canonical and non-canonical Wnt signaling and binds to various Wnt ligands and Frizzled receptors.^{1,4} GPC3-targeting antibodies induce antibody-dependent cellular cytotoxicity (ADCC) and reduce tumor growth in hepatocellular carcinoma (HCC) mouse xenograft models.⁵ Hepatic levels of GPC3 are increased in patients with HCC.⁶ Mutations in *GPC3* are associated with Simpson-Golabi-Behmel syndrome (SGBS), an X-linked condition characterized by pre- and post-natal overgrowth.² Cayman's Glypican-3 Rabbit Monoclonal Antibody (APC) (Clone 024) is composed of a GPC3 monoclonal antibody conjugated to allophycocyanin (APC) and can be used for flow cytometry (FC).

References

- 1. De Cat, B., Muyldermans, S.-Y., Coomans, C., et al. Processing by proprotein convertases is required for glypican-3 modulation of cell survival, Wnt signaling, and gastrulation movements. J. Cell. Biol. 163(3), 625-635 (2003).
- 2. Pilia, G., Hughes-Benzie, R.M., MacKenzie, A., et al. Mutations in *GPC3*, a glypican gene, cause the Simpson-Golabi-Behmel overgrowth syndrome. *Nat. Genet.* **12(3)**, 241-247 (1996).
- 3. Baumhoer, D., Tornillo, L., Stadlmann, S., et al. Glypican 3 expression in human nonneoplastic, preneoplastic, and neoplastic tissues: A tissue microarray analysis of 4,387 tissue samples. Am. J. Clin. Pathol. 129(6), 899-906 (2008).
- 4. Capurro, M., Martin, T., Shi, W., et al. Glypican-3 binds to frizzled and plays a direct role in the stimulation of canonical Wnt signaling. J. Cell. Sci. 127 (Pt. 7), 1565-1575 (2014).
- 5. Ishiguro, T., Sugimoto, M., Kinoshita, Y., et al. Anti-glypican 3 antibody as a potential antitumor agent for human liver cancer. *Cancer Res.* **68(23)**, 9832-9838 (2008).
- 6. Capurro, M., Wanless, I.R., Sherman, M., et al. Glypican-3: A novel serum and histochemical marker for hepatocellular carcinoma. *Gastroenterology* **125(1)**, 89-97 (2003).

ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897