

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Kresoxim-methyl-d-

Item No. 39603

Formal Name: αE-(methoxyimino)-2-[(2-(methyl-d₃)

phenoxy-3,4,5,6-d₄)methyl]-

benzeneacetic acid, methyl ester

MF: $C_{18}H_{12}D_7NO_4$

FW: 320.4


Chemical Purity: ≥95% (Kresoxim-methyl)

Deuterium

Incorporation: \geq 99% deuterated forms (d₁-d₇); \leq 1% d₀

Supplied as: A solid Storage: -20°C Stability: ≥4 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

Kresoxim-methyl-d₇ is intended for use as an internal standard for the quantification of kresoxim-methyl (Item No. 25816) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

Kresoxim-methyl-d₇ is supplied as a solid. A stock solution may be made by dissolving the kresoximmethyl- d_7 in the solvent of choice, which should be purged with an inert gas. Kresoxim-methyl- d_7 is slightly soluble in chloroform and methanol.

Description

Kresoxim-methyl is a strobilurin fungicide. It inhibits conidial germination of V. inaequalis isolates from apple orchards (EC $_{50}$ s = 0.00033-0.0078 mg/L). Kresoxim-methyl also inhibits mycelial growth (EC $_{50}$ = 0.240 mg/L) and is fungicidal against Saprolegnia (MIC = 1 mg/L).2 It increases intracellular calcium levels and disrupts the mitochondrial membrane potential in mouse cortical cultures in a concentration-dependent manner.³ Kresoxim-methyl is toxic to goldfish (C. auratus; LC₅₀ = 0.807 mg/L).² Formulations containing kresoxim-methyl have been used in the control of fungi in agriculture.

References

- 1. Fiaccadori, R., Cicognani, E., Alberoni, G., et al. Sensitivity to strobilurin fungicides of Italian Venturia inaequalis populations with different origin and scab control. Pest Manag. Sci. 67(5), 535-540 (2011).
- Hu, X.-G., Liu, L., Hu, K., et al. In vitro screening of fungicidal chemicals for antifungal activity against Saprolegnia. J. World Aquac. Soc. 44(4), 528-535 (2013).
- Regueiro, J., Olguín, N., Simal-Gándara, J., et al. Toxicity evaluation of new agricultural fungicides in primary cultured cortical neurons. Environ. Res. 140, 37-44 (2015).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information Buyer agrees to purchase the material can be found on our website.

Copyright Cayman Chemical Company, 09/18/2023

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM