

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



## Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



# PRODUCT INFORMATION



### elF2α Activator 2

Item No. 40130

CAS Registry No.: 1501618-04-7

N-[trans-4-[4-(trifluoromethyl)phenoxy] Formal Name:


cyclohexyl]-N'-[3-(trifluoromethyl)phenyl]-urea

Synonym: Eukaryotic Translation Initiation Factor 2

Subunit 1 Activator 2

 $C_{21}H_{20}F_6N_2O_2$ MF:

FW: 446.4 **Purity:** ≥98% Supplied as: A solid Storage: -20°C Stability: ≥4 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

#### **Laboratory Procedures**

elF2 $\alpha$  activator 2 is supplied as a solid. A stock solution may be made by dissolving the elF2 $\alpha$  activator 2 in the solvent of choice, which should be purged with an inert gas.  $elF2\alpha$  activator 2 is soluble in acetonitrile.

#### Description

elF2α activator 2 is an activator of eukaryotic translation initiation factor 2α kinase 1 (ElF2AK1), also known as heme-regulated inhibitor kinase (HRI).<sup>1</sup> It activates HRI and induces eIF2α phosphorylation in a surrogate cell-based reporter assay. eIF2a activator 2 selectively inhibits the proliferation of CRL-2813 melanoma cells (IC<sub>50</sub> = 0.46  $\mu$ M) over non-cancerous NIH3T3 fibroblasts (IC<sub>50</sub> = 3.4  $\mu$ M). It also selectively inhibits the proliferation of CRL-2813 cells transiently transfected with non-targeted siRNA (IC<sub>50</sub> = 0.66  $\mu$ M) over those transiently transfected with HRI-targeted siRNA (IC<sub>50</sub> = 5  $\mu$ M). eIF2 $\alpha$  activator 2 (5  $\mu$ M) increases mRNA expression and protein levels of the elF2α target DNA damage-inducible transcript 3 (DDIT3), also known as C/EBP homologous protein (CHOP), in CRL-2813 cells.<sup>2</sup>

#### References

- 1. Chen, T., Takrouri, K., Hee-Hwang, S., et al. Explorations of substituted urea functionality for the discovery of new activators of the heme-regulated inhibitor kinase. J. Med. Chem. 56(23), 9457-9470 (2013).
- Yefidoff-Freedman, R., Fan, J., Yan, L., et al. Development of 1-((1,4-trans)-4-aryloxycyclohexyl)-3-arylurea activators of heme-regulated inhibitor as selective activators of the eukaryotic initiation factor 2 alpha (EIF2α) phosphorylation arm of the integrated endoplasmic reticulum stress response. J. Med. Chem. 60(13), 5392-5406 (2017).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

## WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information Buyer agrees to purchase the mater can be found on our website.

Copyright Cayman Chemical Company, 12/12/2023

#### **CAYMAN CHEMICAL**

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM