



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



PRODUCT INFORMATION



5,6,7-Trimethoxyflavone

Item No. 40289

CAS Registry No.: 973-67-1

Formal Name: 5,6,7-trimethoxy-2-phenyl-4H-1-benzopyran-4-one

Synonym: Baicalein 5,6,7-trimethyl ether

MF: C₁₈H₁₆O₅

FW: 312.3


Purity: ≥95%

Supplied as: A crystalline solid

Storage: -20°C

Stability: ≥4 years

Item Origin: Synthetic



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

5,6,7-Trimethoxyflavone is supplied as a crystalline solid. A stock solution may be made by dissolving the 5,6,7-trimethoxyflavone in the solvent of choice, which should be purged with an inert gas. 5,6,7-Trimethoxyflavone is soluble in acetonitrile.

Description

5,6,7-Trimethoxyflavone is a flavonoid that has been found in *S. baicalensis* and has diverse biological activities.¹⁻³ It reduces the viability of HepG2 and Hep3B hepatocellular carcinoma cells when used at a concentration of 50 μM.¹ 5,6,7-Trimethoxyflavone decreases herpes simplex virus 1 (HSV-1) and poliovirus replication in HSV-1- or poliovirus-infected Vero cells, as well as decreases cytomegalovirus (CMV) replication in CMV-infected MRC-5 cells (IC₅₀s = 3.2, 32, and 8 mg/L, respectively).² It inhibits LPS-induced nitric oxide (NO), prostaglandin E₂ (PGE₂; Item No. 14010), Tnf-α, IL-1β, and IL-6 production in RAW 264.7 macrophages when used at concentrations of 20 or 40 μM.³ In vivo, 5,6,7-trimethoxyflavone (40 mg/kg) increases survival in a mouse model of LPS-induced sepsis.

References

1. Liao, H.-L. and Hu, M.-K. Synthesis and anticancer activities of 5,6,7-trimethylbaicalein derivatives. *Chem. Pharm. Bull. (Tokyo)* **52**(10), 1162-1165 (2004).
2. Hayashi, K., Hayashi, T., Otsuka, H., et al. Antiviral activity of 5,6,7-trimethoxyflavone and its potentiation of the antiherpes activity of acyclovir. *J. Antimicrob. Chemother.* **39**(6), 821-824 (1997).
3. Rim, H.-K., Yun, C.H., Shin, J.-S., et al. 5,6,7-trimethoxyflavone suppresses pro-inflammatory mediators in lipopolysaccharide-induced RAW 264.7 macrophages and protects mice from lethal endotoxin shock. *Food. Chem. Toxicol.* **62**, 847-855 (2013).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 03/07/2024

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 - USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM