



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



PRODUCT INFORMATION

Iprindole

Item No. 41664

CAS Registry No.: 5560-72-5

Formal Name: 6,7,8,9,10,11-hexahydro-N,N-dimethyl-5H-cyclooct[b]indole-5-propanamine

Synonym: NSC 169449

MF: C₁₉H₂₈N₂


FW: 284.4

Purity: ≥95%

Supplied as: A solution in ethanol

Storage: -20°C

Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Iprindole is supplied as a solution in ethanol. To change the solvent, simply evaporate the ethanol under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as acetonitrile and DMSO purged with an inert gas can be used. Iprindole is sparingly soluble (1-10 mg/ml) in DMSO and slightly soluble (0.1-1 mg/ml) in acetonitrile.

Description

Iprindole is a tricyclic antidepressant.^{1,2} It binds to the serotonin (5-HT) receptor subtypes 5-HT_{2A} and 5-HT_{2C} in radioligand binding assays (K_i s = 0.217 and 0.206 μM, respectively) but only weakly inhibits norepinephrine (NE) and 5-HT uptake in mouse brain slices (IC_{50} s = 9 and 50 μM, respectively).^{1,3} Iprindole reduces immobility in the forced swim test in rats when administered at a dose of 10 mg/kg once per day for 21 days.²

References

- Ross, S.B., Renyi, A.L., and Ogren, S.O. A comparison of the inhibitory activities of iprindole and imipramine on the uptake of 5-hydroxytryptamine and noradrenaline in brain slices. *Life Sci.* **10**(22), 1267-1277 (1971).
- Berettera, C., Invernizzi, R.W., Pulvirenti, L., et al. Chronic treatment with iprindole reduces immobility of rats in the behavioural 'despair' test by activating dopaminergic mechanisms in the brain. *J. Pharm. Pharmacol.* **38**(4), 313-315 (1986).
- Pälvinäki, E.P., Roth, B.L., Majasuo, H., et al. Interactions of selective serotonin reuptake inhibitors with the serotonin 5-HT_{2C} receptor. *Psychopharmacology (Berl)* **126**(3), 234-240 (1996).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.