

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



CCN3 (human, recombinant)

Item No. 41888

Overview and Properties

Synonyms:	Cellular Communication Network Factor 3, IBP-9, IGFBP-9, IGFBP9, IGF-binding Protein 9, Insulin-like Growth Factor-binding Protein 9, NOV
Source:	Active recombinant human C-terminal His-taggged CCN3 expressed in insect cells
Amino Acids:	1-357
Uniprot No.:	P48745
Molecular Weight:	37 kDa
Storage:	-80°C (as supplied)
Stability:	≥1 year
Purity:	≥94% estimated by SDS-PAGE
Supplied in:	Lyophilized from sterile 50 mM Tris, pH 7.4, with 100 mM sodium chloride, 0.5 mM
	phenylmethylsulfonyl fluoride, 10 mM imidazole, and 10% glycerol
Endotoxin Testing:	< 1.0 EU/ μ g, determined by the LAL endotoxin assay
Protein	
Concentration:	<i>batch specific</i> mg/ml
Activity:	<i>batch specific</i> U/ml
Specific Activity:	<i>batch specific</i> U/mg
Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.	

Image



Lane 2: CCN3 (human, recombinant)

SDS-PAGE Analysis of CCN3 (human, recombinant). This protein has a calculated molecular weight of 37 kDa. It has an apparent molecular weight of approximately 47 kDa by SDS-PAGE under reducing conditions due to gycosylation.

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 08/02/2024

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM

PRODUCT INFORMATION



Description

Cellular communication network factor 3 (CCN3) is a growth factor and member of the CCN protein family with roles in stem cell regulation, hematopoiesis, and the regulation of bone growth, as well as cell growth, differentiation, and migration.^{1,2} It is a secreted protein that is associated with the extracellular matrix (ECM) and is composed of a signal peptide linked to four structural domains; an insulin-like growth factor binding protein domain (IGFBP), a von Willebrand type C repeat (VWC), and a thrombospondin type I domain (TSP-1), all associated with protein binding, and a cysteine knot carboxyl terminal that promotes oligomerization. CCN3 is present in various stem cell lineages, including chondrogenic, osteogenic, and hematopoietic lineages, and acts through the core stem cell signaling pathways mediated by Notch and bone morphogenetic protein (BMP).^{1,3} Ccn3 expression is upregulated during early phase bone regeneration, and $Ccn3^{-/-}$ mice exhibit accelerated bone regeneration compared to wild-type mice, indicating Ccn3 is a negative regulator of this process.³ However, Ccn3 expression increases in KiSS1 neurons in the arcuate nucleus (ARCKiSS1) in female mice during lactation, and reduction of Ccn3 expression in ARCKiSS1 induces bone loss in female lactating mice fed a low-calcium diet.⁴ Cayman's CCN3 (human, recombinant) protein can be used for binding assays. This protein consists of 337 amino acids, has a calculated molecular weight of 37 kDa, and a predicted N-terminus of Thr32 after signal peptide cleavage. By SDS-PAGE under reducing conditions, the apparent molecular mass of the protein is 47 kDa due to glycosylation.

References

- 1. McCallum, L. and Irvine, A.E. CCN3 a key regulator of the hematopoietic compartment. *Blood Rev.* **23(2)**, 79-85 (2009).
- Peng, L., Wei, Y., Shao, Y., et al. The emerging roles of CCN3 protein in immune-related diseases. Mediators Inflamm. 2021, 5576059 (2021).
- 3. Matsushita, Y., Sakamoto, K., Tamamura, Y., *et al.* CCN3 protein participates in bone regeneration as an inhibitory factor. *J. Biol. Chem.* **288(27)**, 19973-19985 (2013).
- 4. Babey, M.E., Krause, W.C., Chen, K., et al. A maternal brain hormone that builds bone. Nature (2024).

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM