

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Sphingosine-1-phosphate (d18:1)

Item No. 62570

CAS Registry No.: 26993-30-6

Formal Name: 2S-amino-4E-octadecene-1,3R-diol

1-(dihydrogen phosphate)

S1P (d18:1), Sphingosine-1-Phosphoric Acid Synonyms:

 $C_{18}H_{38}NO_5P$ MF:

379.5 FW: **Purity:** ≥98%

Supplied as: A crystalline solid

Storage: -20°C Stability: ≥4 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Sphingosine-1-phosphate (S1P) (d18:1) is supplied as a crystalline solid. S1P (d18:1) is sparingly soluble in organic solvents. Aqueous solutions can be prepared by directly dissolving the crystalline solid in basic buffers. The solubility of S1P (d18:1) in 0.3 M NaOH is approximately 4 mg/ml, but it will precipitate at pH less than 7.5. The reconstituted material will be stable for 24 hours if stored at 4°C.

Description

Sphingosine-1-phosphate (S1P) is the product of phosphorylation of sphingosine by sphingosine kinase that is secreted from cells and acts as an agonist at S1P receptors. 1,2 It increases intracellular calcium levels in TAg-Jurkat cells expressing ${\rm S1P}_1$ and ${\rm G}_{\rm qi5}$, which allows for phospholipase C stimulation by ${\rm G}_{\rm i}$ proteins, when used at a concentration of 200 nM, as well as in TAg-Jurkat cells expressing S1P₂ and S1P₃ receptors (EC₅₀s = 8 and 11 nM, respectively).³ Intra- and extracellular levels of S1P vary spatially allowing it to function as an autocrine or paracrine factor, respectively, and dysregulation of S1P levels are associated with various disease states, such as inflammation and autoimmunity. S1P has a wide variety of effects, including an involvement in cell growth, angiogenesis, immunity, and neuroprotection.

References

- 1. Sanchez, T. and Hla, T. Structural and functional characteristics of S1P receptors. J. Cell. Biochem. 92(5), 913-922 (2004).
- 2. van Koppen, C.J., zu Heringdorf, D.M., Laser, K.T., et al. Activation of a high affinity G_i protein-coupled plasma membrane receptor by sphingosine-1-phosphate. J. Biol. Chem. 271(4), 2082-2087 (1996).
- 3. An, S., Bleu, T., and Zheng, Y. Transduction of intracellular calcium signals through G protein-mediated activation of phospholipase C by recombinant sphingosine 1-phosphate receptors. Mol. Pharmacol. 55(5), 787-794 (1999).
- 4. Cartier, A. and Hla, T. Sphingosine 1-phosphate: Lipid signaling in pathology and therapy. Science 366(6463), (2019).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 03/28/2024

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM