

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



CPHPC

Item No. 75500

CAS Registry No.: 224624-80-0

Formal Name: 1,1'-(1,6-dioxo-1,6-hexadiyl)bis-D-proline

Synonym: GSK2315698, Miridesap, Ro 63-8695

MF: $C_{16}H_{24}N_2O_6$ FW: 340.4 **Purity:** ≥95%

Supplied as: A crystalline solid

Storage: -20°C Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis

0



CPHPC is supplied as a crystalline solid. A stock solution may be made by dissolving the CPHPC in the solvent of choice. CPHPC is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide, which should be purged with an inert gas. The solubility of CPHPC in these solvents is approximately 30, 20, and 25 mg/ml, respectively.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of CPHPC can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of CPHPC in PBS (pH 7.2) is approximately 10 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

CPHPC is an inhibitor of the protein-protein interaction between serum amyloid P (SAP) and amyloid fibrils $(IC_{50} = 0.9 \mu M)$. It also crosslinks with pentameric SAP in the plasma to form a decameric complex, which is eliminated via hepatic clearance. As plasma and amyloid deposits of SAP exist in a dynamic equilibrium, CPHPC induces the dissociation of SAP from tissue amyloid deposits, leading to its redistribution into the plasma, where it is eliminated. DPHC (1 mg/ml in the drinking water) reduces serum human SAP levels in a transgenic mouse model of amyloidosis.

Reference

1. Pepys, M.B., Herbert, J., Hutchinson, W.L., et al. Targeted pharmacological depletion of serum amyloid P component for treatment of human amyloidosis. Nature 417, 254-259 (2002).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 02/01/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM