

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Sclerotiorin

Item No. 89460

CAS Registry No.: 549-23-5

Formal Name: 7-(acetyloxy)-5-chloro-3-[(1E,3E,5S)-3,5-

dimethyl-1,3-heptadien-1-yl]-7-methyl-

6H-2-benzopyran-6,8(7H)-dione

MF: C₂₁H₂₃CIO₅ 390.9 FW: **Purity:** ≥98%

 λ_{max} : 206, 287, 365 nm UV/Vis.: Supplied as: A crystalline solid

Storage: -20°C Stability: ≥4 years

Item Origin: Fungus/Penicillium sp.

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Sclerotiorin is supplied as a crystalline solid. A stock solution may be made by dissolving the sclerotiorin in an organic solvent purged with an inert gas. Solvents such as ethanol, DMSO, and dimethyl formamide purged with an inert gas can be used. The solubility of sclerotiorin in these solvents is approximately 3, 20, and 30 mg/ml respectively.

Sclerotiorin is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, sclerotiorin should first be dissolved in DMF and then diluted with the aqueous buffer of choice. Sclerotiorin has a solubility of approximately 0.3 mg/ml in a 1:2 solution of DMF:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

Sclerotiorin is a natural product isolated primarily from Penicillium species. It inhibits soybean lipoxygenase-1 with an IC₅₀ value of 4.2 μ M.¹ Sclerotiorin also exhibits a number of other activities including inhibition of cholesterol ester transfer protein (IC_{50} = 19.4 μ M), 2 inhibition of Grb2-Shc interaction (IC_{50} = 22 μ M), 3 and antagonism of endothelin receptors (IC_{50} = 114 and 152 μ M for human ET_A and ET_B, respectively). 4

References

- 1. Chidananda, C. and Sattur, A.P. Sclerotiorin, a novel inhibitor of lipoxygenase from Penicillium frequentans. J. Agric. Food Chem. 55(8), 2879-2883 (2007).
- Tomoda, H., Matsushima, C., Tabata, N., et al. Structure-specific inhibition of cholesteryl ester transfer protein by azaphilones. J. Antibiotics 52(2), 160-170 (1999).
- Ji-Youn, N., Son, K.-H., Kim, H.-K., et al. Sclerotiorin and isochromophilone IV: Inhibitors of Grb2-Shc interaction, isolated from Penicillium multicolor F1753. J. Microbiol. Biotechnol. 10(4), 544-546 (2000).
- 4. Pairet, L., Wrigley, S.K., Chetland, I., et al. Azaphilones with endothelin receptor binding activity produced by Penicillium sclerotiorum: Taxonomy, fermentation, isolation, structure elucidation and biological activity. J. Antibiotics 48(9), 913-923 (1995).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 08/25/2023

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM