

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com


PRODUCT INFORMATION



Thiodiglycolic Acid-d₄

Item No. 9002478

CAS Registry No.: Formal Name:	132090-51-8 2,2'-thio <i>bis</i> (acetic-2,2,2',2'-d₄ acid)
Synonyms:	Dicarboxydimethyl sulfide-d ₄ , Mercaptodiacetic
	Acid-d ₄ , TDGA-d ₄ , Thiodiacetic Acid-d ₄
MF:	C ₄ H ₂ D ₄ O ₄ S
FW:	154.2
Chemcial Purity:	≥98% Thiodiglycolic Acid
Deuterium	
Incorporation:	≥99% deuterated forms (d ₁ -d ₄); ≤1% d ₀
Stability:	≥2 years at -20°C
Supplied as:	A crystalline solid



Laboratory Procedures

Thiodiglycolic acid-d₄ (TDGA-d₄) contains four deuterium atoms at the 2, 2, 2', and 2' positions. It is intended for use as an internal standard for the quantification of TDGA by GC- or LC-mass spectrometry (MS). For long term storage, we suggest that TDGA- d_1 be stored as supplied at -20°C. It should be stable for at least two years.

TDGA- d_4 is supplied as a crystalline solid. A stock solution may be made by dissolving the TDGA- d_4 in the solvent of choice. TDGA-d₄ is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF), which should be purged with an inert gas. The solubility of TDGA-d₄ in ethanol and DMF is approximately 30 mg/ml and approximately 25 mg/ml in DMSO.

TDGA-d₄ is used as an internal standard for the quantification of TDGA by stable isotope dilution MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

Description

TDGA is a product of interaction of the cysteine component of glutathione with two-carbon units released during catabolism. It is found in concentrations below 20 mg/l in urine of healthy individuals and elevated following ingestion of certain drugs.¹ Determination of TDGA concentration in urine has been used to characterize the metabolism of substances participating in methionine synthesis in order to identify imbalances leading to hyperhomocystinuria.² Additionally, because TDGA is used as raw material in the polymer industry, its detection in human urine has served as a biomarker for exposure to carcinogenic vapors such as vinylchloride monomer produced during polymer manufacture.²

References

- 1. Navratil, T., Petr, M., Senholdova, Z., et al. Physiol. Res. 56(1), 113-122 (2007).
- 2. Cheng, Y.-J., Huang, Y.-F., and Ma, Y.-C. J. Occup. Environ. Med. 43(11), 934-938 (2001).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

al should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 11/18/2015

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM