



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TECHNICALLY *Speaking*

Place your order with CEDARLANE® or your local distributor.

Please contact CEDARLANE® for lot specific information.

HEMO-LO GUINEA PIG COMPLEMENT

CL4555

DESCRIPTION:

Cedarlane Hemo-Lo Guinea Pig Complement is designed specifically for sensitive antibody forming cell plaque assays utilizing free or haptenated SRBC as indicator cells. This reagent is pre-absorbed with sheep erythrocytes to reduce background hemolysis. Each batch is thoroughly tested for high complement activity and low background hemolysis, and can be used without further processing.

PRESENTATION: 5 ml, lyophilized

STORAGE: Store at -20°C before reconstitution.

RECONSTITUTION:

Reconstitute with 5 ml of ice cold distilled water. Keep on ice until used. Serum should be used within 2 hours after reconstitution. If smaller volumes are required, aliquot in appropriate volume immediately after reconstitution, freeze immediately, and store frozen material at -70°C. After thawing frozen aliquots, do not refreeze.

NOTE:

1. The reconstituted material will appear cloudy. This is a result of the lyophilization process and does not affect activity.
2. Optimum concentration of this batch of Cedarlane Hemo-Lo Guinea Pig Complement will depend on the system in which it is used. This reagent should be titrated in your system to determine the concentration that provides optimal activity in the absence of background hemolysis.

STERILITY:

This reagent is not sold as sterile, but can be sterilized by filtration if necessary.

**FOR LABORATORY RESEARCH ONLY
NOT FOR HUMAN DRUG OR DIAGNOSTIC USE**

rev/lh/nov/98

For more information or to place an order please contact...

CEDARLANE®
LABORATORIES LIMITED



toll free: 1-800-268-5058
in North America

phone: (905) 878-8891 • fax: (905) 878-7800

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information

website: www.cedarlanelabs.com • e-mail: info@cedarlanelabs.com