



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# TECHNICALLY *Speaking*

Place your order with CEDARLANE® or your local distributor.

Please contact CEDARLANE® for lot specific information.

## Lympholyte®-Rat

CL5040, CL5041, CL5045

### DESCRIPTION:

Lympholyte®-Rat is a density separation medium specifically designed for the isolation of viable lymphocytes from rat lymphoid cell suspensions.

### APPLICATIONS:

Lympholyte®-Rat can be utilized with a simple protocol for the elimination of erythrocytes, dead cells and debris from rat spleen, lymph node, thymus and bone marrow suspensions. The resulting cell population demonstrates a high and non-selective recovery of viable lymphocytes, which are suitable for use as target cells in cytotoxicity and FACS assays, as well as in *in vivo* and *in vitro* functional studies. Other successful applications include:

- i) the removal of dead cells in sequential cytotoxicity studies eg. B-cell depletion.
- ii) the removal of erythrocytes, dead cells and debris from other rat tissue suspensions including liver and lung.
- iii) the harvesting of viable cells and removal of dead cells and debris from various clone cell and hybridoma cell lines.

### PRESENTATION:

Sterile liquid.  
CL5040, 5 x 30 ml  
CL5045, 1 x 500 ml  
CL5041 1x 100 ml

### STORAGE/STABILITY:

Store at room temperature unopened. Store at +4°C once opened. *Always store protected from light.*

Note: Phase separation may occur with long-term storage. **SHAKE WELL BEFORE USE. ALLOW TO STAND UNTIL NO AIR BUBBLES PRESENT. USE AT ROOM TEMPERATURE.**

*Continued Overleaf...*

For more information or to place an order please contact...

**CEDARLANE®**  
**LABORATORIES LIMITED**



*toll free: 1-800-268-5058*  
*in North America*

phone: (905) 878-8891 • fax: (905) 878-7800

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information  
website: [www.cedarlanelabs.com](http://www.cedarlanelabs.com) • e-mail: [info@cedarlanelabs.com](mailto:info@cedarlanelabs.com)

## **SPECIFICATIONS:**

**Composition:** Nycograde™ Polysucrose 400 and Sodium Diatrizoate

**Density:**  $1.094 \pm 0.001 \text{ g/cm}^3$  @ 22°C.

**pH:**  $6.9 \pm 0.3$

**Viability/ Purity:** Recovery of viable lymphocytes  $\geq 70\%$ .

Results obtained on a rat spleen suspension:

<b>Fraction</b>	<b>Viable Lymphocytes</b>	<b>Erythrocyte Contamination</b>
upper	<1%	0
interphase	>70%	< 15%
lower	<10%	< 5%
pellet	<20%	> 80%

## **METHOD OF USE:**

Use Lympholyte®-Rat and preferably a serum-free medium of choice (Phosphate Buffered Saline, Modified McCoy's Medium, etc.) at room temperature (approximately 22°C).

1. Prepare a lymphocyte suspension using your preferred method and medium. Spleen has a high membrane content and a clean suspension is required for proper separation.

Suggested method: a) cut up spleen into small pieces  
b) homogenize  
c) pass suspension through a fine screen mesh

Other tissues: homogenize thoroughly to obtain a clean suspension.

2. Adjust the cell concentration to  $2 \times 10^7$  nucleated cells per ml or less.  
**Note:** .If cell suspension contains a large amount of debris or erythrocytes, a cleaner separation will be obtained if the cell concentration is set at  $1.0 \times 10^7$  cells/ml.
3. Layer the cell suspension over Lympholyte®-Rat according to Method A or Method B (see figures). Use a 10-15 ml centrifuge tube.

**Method A:** Add 5 ml of Lympholyte®-Rat to the centrifuge tube. Using a pipette, carefully layer 5 ml of the cell suspension over the Lympholyte®-Rat with as little mixing as possible at the interface (Figure A). Since Lympholyte®-Rat is of greater density than the cell suspension, a distinct interface will be formed (Figure C).


**Method B:** Add 5 ml of the cell suspension to the centrifuge tube. Place a large (23 cm) Pasteur pipette to the bottom of the tube

(Figure B). Slowly add Lympholyte®-Rat to the Pasteur pipette allowing gravity to layer it under the cell suspension. Continue until

5 ml of Lympholyte®-Rat has been layered under the cell suspension. Since Lympholyte®-Rat is of greater density than the cell suspension, the cell suspension will form a layer above the Lympholyte®-Rat with a distinct interface (Figure C).

4. Centrifuge for 20 minutes at 1000-1500g at **room temperature**.
5. After centrifugation, there will be a well-defined lymphocyte layer at the interface (Figure D). Using a Pasteur pipette, carefully remove the cells from the interface and transfer to a new centrifuge tube.
6. Dilute the transferred cells with medium to reduce the density of the solution. Centrifuge at 800g for 10 minutes to pellet the lymphocytes, then discard the supernatant.
7. Wash the lymphocytes 2-3 times in medium (can now use media containing serum) before further processing.

*Continued Overleaf....*



**FOR RESEARCH USE ONLY**

NYCOGRADE™ is a registered trademark of Nycomed Pharma AS, Oslo, Norway.

LYMPHOLYTE® is a registered trademark of Cedarlane Laboratories Ltd., Burlington, Canada.

**References:**

1. Kawashima, H., *et al.* 1994. *Journal of Immunology*. **153**: 1982.