



**SZABO  
SCANDIC**

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](http://linkedin.com/company/szaboscandic)



# Technically Speaking



[www.cedarlanelabs.com](http://www.cedarlanelabs.com)

Conveniently Delivering You Today's Innovations  
for the Science of Tomorrow™

## Anti-Human IgG<sub>3</sub> (Hinge Region) Monoclonal Antibody

Catalogue#	Format	Size	Concentration	Isotype Control
<b>CL6050AP</b>	Purified	50 µg	1.0 mg/ml	CLCMG100
<b>CL6050AP-2</b>	Purified	250 µg	1.0 mg/ml	CLCMG100
<b>CL6050AP-3</b>	Purified	500 µg	1.0 mg/ml	CLCMG100
<b>CL6050B</b>	Biotin	100 µg	0.1 mg/ml	CLCMG115
<b>CL6050F</b>	FITC	100 µg	0.1 mg/ml	CLCMG101
<b>CL6050PE</b>	PE	50 µg	0.1 mg/ml	CLCMG104
<b>CL6050HP</b>	HRPO	100 µg	1.0 mg/ml	CLCMG107

Isotype: Mouse IgG<sub>1</sub>

### DESCRIPTION:

There are four subclasses of IgG present in humans, IgG1, IgG2, IgG3 and IgG4. These IgGs are secreted by B cells and are found in blood and extracellular fluids and provide protection against infections caused by bacteria, fungi and viruses. IgG3 is unique in that it has an extended hinge region of the heavy chain conferring greater relative flexibility of the Fab arms with respect to the Fc region. This greater flexibility allows IgG3 to be particularly effective in the induction of effector functions and inflammatory responses. Its shorter half-life may function to limit the potential of excessive inflammatory responses.

Cedarlane's mouse anti-human IgG3 antibody, clone HP6050 recognizes the hinge region of the heavy chain of human IgG3.

### PRESENTATION:

**Purified:** Purified IgG buffered in PBS and 0.02% NaN<sub>3</sub>. (Purified via Protein G Chromatography). For maximum recovery of contents, spin down tube before use.

**Biotin, FITC, and PE:** Biotin/FITC/PE conjugated IgG buffered in PBS, 0.02% NaN<sub>3</sub> and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/ml.

**HRPO:** HRPO conjugated IgG buffered in PBS with 40% glycerol (v/v) and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/ml. No NaN<sub>3</sub> with other preservatives.

### STORAGE/STABILITY:

For all formats, store at + 4°C. DO NOT FREEZE PE conjugates. For long term storage (**Purified/Biotin/FITC/HRPO**), aliquot and freeze unused portion at -20°C in volumes appropriate for single usage. Avoid freeze/thaw cycles. For Maximal recovery of contents, please quick-spin vial before opening.

*Continued Overleaf.....*

Visit our website for your local distributor.



[www.cedarlanelabs.com](http://www.cedarlanelabs.com)

An ISO 9001:2000 and ISO 13485:2003  
registered company.

**In CANADA: Toll Free: 1-800-268-5058**

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020  
e-mail: general@cedarlanelabs.com

**In the USA: Toll Free: 1-800-721-1644**

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138  
e-mail: service@cedarlanelabs.com

## **APPLICATION:**

This antibody is suitable for use in ELISA (both capture and detection antibody), Western Blot, Flow Cytometry and Immunohistochemistry with paraffin-embedded and frozen sections.

**\* For optimal results in various applications, it is recommended that each investigator determine dilutions appropriate for individual use.**

## **SPECIFICATIONS:**

Clone: HP6050

Immunogen: Human IgG<sub>3</sub> myeloma protein.

Specificity: Human IgG<sub>3</sub> Hinge Region.

ELISA Specificity: human IgG subclass 1 – <0.01%  
human IgG subclass 2 – <0.01%  
human IgG subclass 3 – 100%  
human IgG subclass 4 – <0.01%

## **REFERENCES:**

1. Vidarsson G, Dekkers G, Rispens T. IgG subclasses and allotypes: from structure to effector functions. *Front Immunol.* 5:520, 2014.
2. Hamilton, R.G. et al; Isoelectric focusing-affinity immunoblot analysis of mouse monoclonal antibodies to the four human IgG subclasses. *Electrophoresis*, 8:127-34, 1987.
3. Papadea, C. et al; Monoclonal antibody-based solid-phase immunoenzymometric assays for quantifying human immunoglobulin G and its subclasses in serum. *Clin. Chem.*, 31: 1940-5, 1985.
4. Jeffries, R. et al; Evaluation of monoclonal antibodies having specificity for human IgG sub-classes: results of an IUIS/WHO collaborative study; *Immunol Lett.* 10(3-4):223-52, 1985.
5. Reimer, C.B. et al; Evaluation of thirty-one mouse monoclonal antibodies to human IgG epitopes. *Hybridoma*, 3: 263-75, 1984.

**FOR RESEARCH USE ONLY**

JK 07/09/2018