

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





for the Science of Tomorrow™

Anti-Mouse CD3z (CD247) **Monoclonal Antibody**

Catalogue#	Format	Size	Concentration	Isotype Control
CL7230AP	Purified	250 μg	1.0 mg/ml	CLCHM00
CL7230B/-3	Biotin	100/300 μg	0.1 mg/ml	CLCHM15
CL7230F/-3	FITC	100/300 μg	0.1 mg/ml	CLCHM01

Isotype: Hamster IgG2κ

DESCRIPTION:

Cedarlane's Purified Anti-Mouse CD3z Monoclonal Antibody detects the zeta subunit of the T cell receptor (TCR). The TCR complex consists of a ligand specific alpha/beta heterodimer non-covalently associated with five invariant chains including the CD3 gamma, delta, eta and zeta subunits. This complex regulates assembly and expression of the receptor and is thought to be responsible for transmembrane transduction of signals after binding to $TCR\alpha\beta$.

Studies have suggested that the CD3z subunit (also known as CD247) plays an important role in two distinct cellular compartments. In the cytoplasm, it may function to regulate Ag receptor expression and in the plasma membrane it may be required for optimal signaling by the physiologic ligand Ag/MHC. The CD3z subunit is also associated with the CD16 IgG Fc receptor on NK cells.

This clone (H146-968) has been reported to work in flow cytometry (in permeabilized cells), Western Blotting and ELISA.

PRESENTATION:

Purified: Purified IgG buffered in PBS and 0.02% NaN3. (Purified from ascitic fluid via Protein G Chromatography).

Biotin and FITC: Biotin/FITC conjugated IgG buffered in PBS, 0.02% NaN₃ and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/mL.

STORAGE/STABILITY:

For all formats, store at 4°C. DO NOT FREEZE PE conjugates. For long term storage (Purified and Biotin), aliquot and freeze unused portion at -20°C in volumes appropriate for single usage. Avoid freeze/thaw cycles.

Visit our website for your local distributor.



In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020 e-mail: general@cedarlanelabs.com

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138 e-mail: service@cedarlanelabs.com

SPECIFICATIONS:

Clone: H146-968

Hybridoma Production:


Immunization:

Immunogen: KLH conjugated mouse zeta-chain (CD3) sequence 151-164

Donor: Armenian hamster

Specificity: Mouse CD3z (zeta chain); also known as CD247

TEST RESULTS:



C57/BL6 mouse thymocytes (fixed and permeabilized) were stained with anti-CD3z (clone: H146-968) (filled histogram) or Armenian hamster IgG isotype control (open histogram).

N.B. Appropriate control samples should always be included in any labeling studies.

* For optimal results in various applications, it is recommended that each investigator determine dilutions appropriate for individual use.

REFERENCES:

- 1. Rozdzial, MM, Kudo RT, Turner SL and Finkel TH. 1994. Developmental Regulation of the TCR ζ -Chain. J Immunol 153:1563.
- 2. Moingeon P *et al.* 1992. CD3ζ Dependance of the CD2 Pathway of Activation in T Lymphocytes and Natural Killer Cells. *Proc Natl Acad Sci USA*. February 15; 89 (4):1492.