

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



## Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



# Agrisera

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

#### Product no AS16 4083

#### HXK1 | Hexokinase 1 (Chlamydomonas)

#### **Product information**

Immunogen KLH-conjugated synthetic peptide derived from Chlamydomonas reinhardtii hexokinase-1, UniProt: A8JGU7

**Host** Rabbit

Clonality Polyclonal

Purity Serum

Format Lyophilized

**Quantity** 50 μ

**Reconstitution** For reconstitution add 50 μl of sterile water.

Storage Store lyophilized/reconstituted at -20°C; once rec

Store lyophilized/reconstituted at -20 °C; once reconstituted make aliquots to avoid repeated freeze-thaw cycles. Please, remember to spin tubes briefly prior to opening them to avoid any losses that might occur from lyophilized

material adhering to the cap or sides of the tubes.

#### **Application information**

Recommended dilution 1:1000 (WB)

Expected | apparent

w

67.3 kDa

**Confirmed reactivity** 

Chlamydomonas reinhardtii, Eudorina elegans, Gonium pectorale


**Predicted reactivity** 

Species of your interest not listed? Contact us

Selected references

<u>Upadhyaya</u> and Jagadeeshwar Rao (2019). Reciprocal regulation of photosynthesis and mitochondrial respiration by TOR kinase in Chlamydomonas reinhardtii. Plant Direct Volume 3, Issue 11.

#### Application example



Total protein of *Chlamydomonas reinhardtii* strain 21gr (1), strain cw15 (2), strain M10 (3) was extracted using 1x Laemmli buffer + DTT + protease and phosphatase inhibitors without bromophenol blue heated at 95 °C for ten minutes then vortexed with zirconia beads at max speed for ten minutes. Lysate was spun at 4 °C for ten minutes at max speed in a tabletop centrifuge. Proteins were precipitated out of supernatant in 80% acetone at -20 °C for 20 min, spun at 4 °C at max speed for 10 min. Pellet was washed 1x with 80% acetone and resuspended in 1x LDS sample buffer + DTT + inhibitors at a concentration of 1 µg chlorophyll/uL. Protein equivalent to 1 of chlorophyll was loaded on a 4-20% Bis-Tris gel and blotted to PVDF using wet transfer methods. Blot was blocked with 10% milk in TBS-T for 1hr/RT. Blot was incubated in the primary antibody at a dilution of 1:1000 in 5% milk in TBS-T ON/4 °C. The antibody solution was decanted and the blot was rinsed 3 times for 5 min in TBS-T at RT with agitation. Blot was incubated in Agrisera matching secondary diluted to 1:50 000 (AS09 602) in 5% milk in TBS-T for 1hr/RT with agitation. The blot was washed as above and developed withchemiluminescent detection reagent and imaged on a Licor C-Digit blot scanner. Exposure time was 12 min.

Courtesy of Jessica Rakijas, Kansas State University, USA