



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



TECHNICALLY *Speaking*

Place your order with CEDARLANE® or your local distributor.

Please contact CEDARLANE® for lot specific information.

Recombinant Human S100P (rHuS100P)

CLPRO425
CLPRO425-2
Lot: 107S100P

Background:

S100P was originally described as a placental protein of 95 amino acid residues shares about 50% sequence identity with the brain S100 proteins alpha and beta. S100 proteins are small dimeric members of the EF-hand superfamily of $\text{Ca}^{(2+)}$ binding proteins thought to participate in mediating intracellular $\text{Ca}^{(2+)}$ signals by binding to and thereby regulating target proteins in a $\text{Ca}^{(2+)}$ -dependent manner. S100P is dysregulated in the androgen-independent prostate cancer cell lines LNCaP-R, DU145, and PC3 and may play a role in the etiology of prostate cancer. In ductal hyperplasias, *in situ* and invasive ductal carcinoma, but not in the normal tissues, S100P overexpression is an early event that might play an important role in the immortalization of human breast epithelial cells in vitro and tumor progression in vivo. In NIH3T3 cells, the expression of S100P led to the presence of S100P in the culture medium, increased cellular proliferation, and enhanced survival following detachment from the culture substrate or after exposure to the chemotherapeutic agent 5-fluorouracil. The proliferation and survival effects of S100P expression were duplicated in a time- and concentration-dependent manner by extracellular addition of purified S100P to wild-type NIH3T3 cells and correlated with the activation of Erks and NFB.

To determine the mechanisms involved in these effects, the hypothesis was tested that S100P activated RAGE (Receptor for Activated Glycation End-Products). It was found that S100P coimmunoprecipitated with RAGE. Furthermore, the effects of S100P on cell signaling, proliferation and survival were blocked by agents that interfere with RAGE including administration of an amphoterin derived peptide known to antagonize RAGE activation, anti-RAGE antibodies and by expression of a dominant negative RAGE. These data suggest that S100P can act in an autocrine manner via RAGE to stimulate cell proliferation and survival.

Description:

The Recombinant Human S100P has a molecular mass of 10.4 kDa containing 95 amino acid residues of the human S100P.

Source: *E.Coli*

Presentation:

2 µg (CLPRO425) or 10 µg (CLPRO425-2), sterile filtered lyophilized (freeze-dried) powder. The protein was lyophilized from 0.5 mg/ml in 0.05 M Phosphate buffer, pH7.2, 0.1 M NaCl.

Continued Overleaf...

For more information or to place an order please contact...

CEDARLANE®
LABORATORIES LIMITED



toll free: 1-800-268-5058
in North America

phone: (905) 878-8891 • fax: (905) 878-7800

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information
website: www.cedarlanelabs.com • e-mail: info@cedarlanelabs.com

Specificity:

The recombinant human S100P is 100% homologous with the human S100P.

Amino Acid sequence:

MTELEAAMGM IIDVFSRYSG SEGSTQTLTK GELKVLMEKE LPGFLQSGKD KDAVDKLLKD
LDANGDAQVD FSEFIVFVAA ITSACHKYFE KAGLK

Purification Method:

Two-step procedure using anion-exchange and size exclusion chromatography.

Purity:

>95% (SDS-PAGE analyzed).

Reconstitution:

Add 0.2 ml of deionized H₂O and let the lyophilized pellet dissolve completely.

Storage:

Store lyophilized protein at -20°C. Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles. Reconstituted protein can be stored at 4°C for a limited period of time; it does not show any change after two weeks at 4°C.

Applications:

Western blotting

References:

- Jin G, Wang S, Hu X, Jing Z, Chen J, Ying K, Xie Y, Mao Y. Characterization of the tissue-specific expression of the s100P gene which encodes an EF-hand Ca²⁺-binding protein. Mol Biol Rep. 2003 Dec; 30(4): 243-8.
- Arumugam T, Simeone DM, Schmidt AM, Logsdon CD. S100P stimulates cell proliferation and survival via RAGE. J Biol Chem. 2003 Nov 14
- Shyu RY, Huang SL, Jiang SY. Retinoic acid increases expression of the calcium-binding protein S100P in human gastric cancer cells. J Biomed Sci. 2003 May-Jun; 10(3): 313-9.
- Sato N, Hitomi J. S100P expression in human esophageal epithelial cells: Human esophageal epithelial cells sequentially produce different S100 proteins in the process of differentiation. Anat Rec. 2002 May 1; 267(1): 60-9.
- Gribenko A, Lopez MM, Richardson JM 3rd, Makhatadze GI. Cloning, overexpression, purification, and spectroscopic characterization of human S100P. Protein Sci. 1998 Jan; 7(1): 211-5.

FOR RESEARCH USE ONLY