



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Human Thyroglobulin

CLPRO443
CLPRO443-2
CLPRO443-3
Lot: 808HTG01

Introduction: Thyroglobulin (TG) represents one of the main autoantigenic targets in autoimmune thyroid disease of humans. TG is a large globular dimeric glycoprotein with a total molecular weight of 660 kDa, which occupies a key precursor role in the biosynthesis of the thyroid hormones. Approximately 75% of the total protein content of the thyroid follicle consists of TG. The thyroid gland uses the Thyroglobulin in order to produce the thyroid hormones thyroxine (T4) and triiodothyronine (T3). Thyroglobulin is produced by the thyroid epithelial cells (thyrocytes) which form spherical follicles. Thyroglobulin is subsequently secreted and stored in the follicular lumen.

Patients with Hashimoto's thyroiditis or Graves' disease, frequently develop antibodies against Thyroglobulin. Tg-specific antibodies help in the diagnosis of the above diseases, however they also may be present in apparently healthy euthyroid individuals. Blood Thyroglobulin levels can be used as a tumor marker for certain kinds of thyroid cancer, and they may also be elevated in cases of Graves' disease.

Description: Human Thyroglobulin is a glycosylated, polypeptide chain having a total molecular mass of 662 kDa.

Synonyms: Thyroglobulin, TGN, AITD3, TG.

Source: Native, Isolated from human thyroid glands.

Presentation: 20 µg (**CLPRO443**), 100 µg (**CLPRO443-2**), or 1 mg (**CLPRO443-3**), human thyroglobulin is supplied at a concentration of 1.1 mg/ml in 16 mM HEPES buffer pH 7.6, 240 mM sodium chloride, and 20% glycerol.

Storage Conditions: Store at 4°C if entire vial will be used within 2-4 weeks.
Store frozen at -20°C for longer periods of time.
Avoid multiple freeze-thaw cycles.

Immunological functions:

1. Binds to IgG-type human auto-antibodies.
2. Auto-antibodies to thyroglobulin recognize conformation-dependent epitopes.
3. Standard ELISA test (checker-board analysis of positive/negative sera panels, including 1st international reference serum NIBSC 65/93).

Coating concentration: 0.7-1 µg/ml (depending on the type of ELISA plate and coating buffer).
Suitable for biotinylation and iodination.

Purity: Greater than 95% as determined by SDS-PAGE.

Laboratory Reagent For Research Use Only

JV 12/15/10

Visit our website for your local distributor.

CEDARLANE®



www.cedarlanelabs.com

An ISO 9001:2000 and ISO 13485:2003
registered company.

In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020
e-mail: general@cedarlanelabs.com

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138
e-mail: service@cedarlanelabs.com