



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Human Fibronectin

CLPRO448
CLPRO448-2
CLPRO448-3

Description: Plasma fibronectin level is elevated in severe coronary artery disease. Increased plasma fibronectin levels are related with venous thromboembolism (VTE) particularly in males, and extend the probable association between biomarkers and risk factors for arterial atherothrombosis and VTE. Fibronectin plays a role in several cellular processes, including tissue repair, embryogenesis, blood clotting, and cell migration/adhesion. Fibronectin consists in two main forms: 1) as an insoluble glycoprotein dimer that serves as a linker in the extracellular matrix and 2) as a soluble disulphide linked dimer found in the plasma. The plasma form is produced by hepatocytes, and the ECM form is synthesized by fibroblasts, chondrocytes, endothelial cells, macrophages, as well as certain epithelial cells. Fibronectin also takes part as a general cell adhesion molecule by anchoring cells to collagen or proteoglycan substrates. Fibronectin organizes cellular interaction with the ECM by binding to different components of the extracellular matrix and to membrane-bound Fibronectin receptors on cell surfaces. Molecular weight: 450 kDa.

Source: Human Plasma

Presentation: 200 µg (CLPRO448), 1 mg (CLPRO448-2), or 10 mg (CLPRO448-3), sterile filtered white lyophilized (freeze-dried) powder. The Fibronectin was lyophilized from a non-sterile 2mg/ml buffer of 10mM sodium phosphate, pH 7.5 and 0.15M NaCl.

Solubility: Purified fibronectin has a tendency to form insoluble aggregates upon reconstitution. We suggest to reconstitute the 1 mg Fibronectin with a chaotropic agent such as urea at room temperature at a concentration of 0.2 mg/ml using sterile water. Let stand 1-2 hours.

Stability: Store the lyophilized fibronectin at 4°C. Upon reconstitution, fibronectin should be stored at 4°C for 2 weeks and for future use below -18°C. Please prevent freeze-thaw cycles.

Purity: Greater than 95.0% as determined by SDS-PAGE.

Cell Culture Application: Fibronectin is useful for the induction of cell attachment to a variety of surfaces including plastic and glass tissue culture labware, petri dishes, coverslips, microcarrier beads, etc. Fibronectin is useful for growth and maintenance of cells in low serum conditions. In general, the reconstituted fibronectin should be diluted with sterile physiological saline or serum-free medium to a concentration of 10-50 µg/ml. The plates are treated with the solution, at room temperature 1-2 hours, the supernatant removed and the plates rinsed with fresh media or buffer. Growth media is then added and the coated plates seeded with a suspension of freshly harvested cells. Care should be taken to inactivate serum proteases used in the preparation of the cells, since fibronectin is very sensitive to proteolysis. Optimum concentration of fibronectin depends on the types of cell and the particular surface used. Generally, use 1-5 µg/cm² of growth surface.

Laboratory Reagent For Research Use Only

MW 11/06/20

Visit our website for your local distributor.

CEDARLANE®



www.cedarlanelabs.com

An ISO 9001:2015 and ISO 13485:2016
registered company.

In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020
e-mail: general@cedarlanelabs.com

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138
e-mail: service@cedarlanelabs.com