



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



TECHNICALLY Speaking

Place your order with CEDARLANE® or your local distributor.

Please contact CEDARLANE® for lot specific information.

Cardiac Actin

CLPRO519

CLPRO519-2

Lot:

Description: Ultra pure Cardiac Actin having a molecular mass of 43,000 daltons.

Presentation: 2 µg (**CLPRO519**) or 10 µg (**CLPRO519-2**) sterile filtered lyophilized powder. The protein was lyophilized from a 1mg/ml solution contains 10mM TRIS/acetate buffer pH 8.0, 0.2 mM CaCl₂, 0.2 mM ATP, 1mM DTT and 0.5% SDS.

Stability: Lyophilized Cardiac-Actin although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Cardiac Actin should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid freeze-thaw cycles.

Solubility: It is recommended to reconstitute the lyophilized Cardiac Actin in sterile 18MΩ-cm H₂O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

Applications:

Protein standard in 1D and 2D SDS gel electrophoresis

Immunoassays

Immunization

Purity:

Greater than 98.0% as determined by RP-HPLC and SDS-PAGE analysis.

References:

1. Transgenic overexpression of cardiac actin in the mouse heart suggests coregulation of cardiac, skeletal and vascular actin expression. *Transgenic Res* 2004 Dec; 13(6):5 31-40
2. A novel complex regulates cardiac actin gene expression through interaction of Emb, a class VI POU domain protein, MEF2D, and the histone transacetylase p300. *Mol Cell Biol* 2004 Apr; 24 (7):2944-57
3. Clinical and genetic characteristics of alpha cardiac actin gene mutations in hypertrophic cardiomyopathy. *J Med Genet* 2004 Jan; 41(1):e10
4. No variants in the cardiac actin gene in Finnish patients with dilated or hypertrophic cardiomyopathy. *Am Heart J* 2002 Jun; 143(6):E6
5. Evaluation of the cardiac actin gene in Doberman Pinschers with dilated cardiomyopathy. *Am J Vet Res* 2001 Jan; 62 (1):33-6
6. Epidemiology of desmin and cardiac actin gene mutations in a european population of dilated cardiomyopathy. *Eur Heart J* 2000 Nov;21 (22):1872-6

FOR RESEARCH USE ONLY

JJ 11/22/06

For more information or to place an order please contact...

CEDARLANE®
LABORATORIES LIMITED

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

toll free: 1-800-268-5058

in North America

phone: (905) 878-8891 • fax: (905) 878-7800

or visit our website for a list of our international distributors including contact information

website: www.cedarlanelabs.com • e-mail: info@cedarlanelabs.com