



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Human Alpha-2 Macroglobulin Protein (A2M)

CLPRO551
CLPRO551-2
CLPRO551-3
Lot: 517PA2M03

Introduction: Alpha-2 Macroglobulin is a serine proteases inhibitor, which inhibits coagulation by inactivating thrombin and Kallikrein, it inhibits fibrinolysis by inactivating plasmin and involved in insulin transport. Alpha-2-Macroglobulin is a large plasma protein, which is produced by the liver, it's composed of 4 identical subunits bound together by -S-S- bonds. A2M is able to inactivate many kinds of proteinases (including serine-, cysteine-, aspartic- and metalloproteinases). α 2M has a 35 amino acid "bait" region in its structure. Proteinases bind and cleave the "bait" region become bound to α 2M. Macrophage receptors recognize the proteinase- α 2M complex and clear it from the system. A2M binds to and removes MMP-2 and MMP-9 (active forms of the gelatinase) from the circulation using scavenger receptors on the phagocytes. The levels of Alpha-2-macroglobulin are increased in nephrotic syndrome which is a condition where the kidneys start to leak out some of the smaller blood proteins. Due to its large size, α 2-macroglobulin is retained in the bloodstream. Increased production of all proteins causes α 2-macroglobulin concentration to increase. Chronic renal failure might lead to amyloid by alpha-2-macroglobulin. A2M is raised in cirrhosis, pregnancy and diabetes.

Description: Human Alpha-2 Macroglobulin is a tetrameric glycoprotein, produced in human plasma and having a molecular mass of 725 kDa.

Synonyms: Alpha-2-macroglobulin, Alpha-2-M, A2M, CPAMD5, FWP007, S863-7, alpha 2M, DKFZp779B086.

Source: Human plasma

Presentation: 200 μ g (CLPRO551), 1 mg (CLPRO551-2), or 10 mg (CLPRO551-3), sterile filtered white lyophilized (freeze-dried) powder. Lyophilized from 0.02 M TRIS, 0.15 M NaCl and 0.1 M sucrose, pH 8.0.

Stability: Human Alpha-2 Macroglobulin although stable at room temperature for 3 weeks, should be stored between 2-8°C.

Purity: Greater than 96.0 %.

Human Virus Test: Serum from each donor has been tested and found negative for antibody to HIV-1, HIV-2, HCV, HBSAG

Laboratory Reagent For Research Use Only

MW 10/31/17

Visit our website for your local distributor.

CEDARLANE®



www.cedarlanelabs.com

An ISO 9001:2000 and ISO 13485:2003
registered company.

In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020
e-mail: general@cedarlanelabs.com

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138
e-mail: service@cedarlanelabs.com