



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Technically
Speaking

CEDARLANE®
www.cedarlanelabs.com



Delivering You Today's Innovations for
the Science of Tomorrow™

Recombinant Mouse Cystatin C

CLPRO597
CLPRO597-2
CLPRO597-3

Lot:

Introduction: Cystatins are a superfamily of cysteine proteinase inhibitors found in both plants and animals. They comprise a group of proteinase inhibitors, widely distributed in tissues and body fluids, and form tight complexes with cysteine proteases such as cathepsin B, H, L and S. Cystatin C, a secreted molecule of this family, is of interest from biochemical, medicine and evolutionary points of view. Cystatin C, with molecular weight of 13260 Da, is composed of 120 amino acids, lacks carbohydrate and has two disulfide bridges located near the carboxyl terminus. Cystatin C is increased in patients with malignant diseases, and is related to the insufficiency of renal function and appears to be a better marker than creatinine. On the other hand, low levels of cystatin C involve cause the breakdown of the elastic laminae and, subsequently, the atherosclerosis and abdominal aortic aneurysm.

Description: Cystatin-C Murine Recombinant produced in *E.Coli* is a single, non-glycosylated, polypeptide chain containing 134 amino acids and having a molecular mass of 15kDa.

The Mouse Cystatin-C is fused to His tag at N-Terminus.

The Mouse Cystatin-C is purified by proprietary chromatographic techniques.

Synonyms: Post G-globulin, CST 3, CST3, Gamma-Trace, Cystatin 3, Amyloid Angiopathy and Cerebral Hemorrhage, Cystatin-C precursor, neuroendocrine basic polypeptide.

Source: *Escherichia Coli*.

Presentation: 2 µg (CLPRO597), 10 µg (CLPRO597-2), or 1 mg (CLPRO597-3), sterile filtered white lyophilized (freeze-dried) powder. The sterile filtered concentrated (0.5 mg/ml) protein solution was lyophilized with 20 mM Tris & 50 mM NaCl pH 7.5.

Solubility: Add 0.1 M Acetate buffer pH 4 to a working concentration of 0.5 mg/ml and let the lyophilized pellet dissolve completely.

Stability: Store lyophilized protein at -20°C. Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles. Reconstituted protein can be stored at 4°C for a limited period of time.

Amino Acid Sequence: MRGSHHHHHH GMASATPKQG PRMLGAPEEA DANEEGVRRR LDFAVSEYNK
GSNDAYHSRA IQVVRARKQL VAGVNYFLDV EMGRTTCTKS QTNLTDCPFH DQPHLMRKAL
CSFQIYSVPW KGTHSLTKFSCKNA

Purity: Greater than 95% as determined by SDS-PAGE.

Laboratory Reagent For Research Use Only

IV 06/25/09

Visit our website for your local distributor.

CEDARLANE®



www.cedarlanelabs.com

An ISO 9001:2000 and ISO 13485:2003
registered company.

In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020
e-mail: general@cedarlanelabs.com

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138
e-mail: service@cedarlanelabs.com