



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Technically
Speaking

CEDARLANE®
www.cedarlanelabs.com



Conveniently Delivering You Today's Innovations
for the Science of Tomorrow™

Recombinant Human B-Cell Leukemia/Lymphoma 2 -NWGR

CLPRO634
CLPRO634-2
CLPRO634-3

Introduction: BCL2 gene encodes an integral outer mitochondrial membrane protein that blocks the apoptotic death of some cells such as lymphocytes. Constitutive expression of BCL2, such as in the case of translocation of BCL2 to Ig heavy chain locus, is thought to be the cause of follicular lymphoma. Two transcript variants, produced by alternate splicing, differ in their C-terminal ends.

Description: Bcl-2 Des NWGR domain (143-146 residues) Human Recombinant produced in *E. Coli* is a single, non-glycosylated polypeptide chain containing 214 amino acids 1-142 and 147-218. The Bcl-2 is expressed as His-Tag fusion protein and purified by proprietary chromatographic techniques.

Synonyms: Apoptosis regulator Bcl-2, BCL2, B-cell CLL/lymphoma 2, Bcl-2.

Source: *Escherichia Coli*.

Presentation: 2 µg (CLPRO634), 10 µg (CLPRO634-2), or 1 mg (CLPRO634-3), sterile filtered white lyophilized (freeze-dried) powder. The protein contains 10mM Tris-HCL pH-8, 1mM EDTA and 250mM NaCl.

Solubility: Suspend Bcl-2 in 100µl of 0.5M Acetic acid, overnight at 4°C. Dilute 10-fold into selected buffer system. BCL-2 has a tendency to form intramolecular disulfide bond, 5mM DTT is recommended in assay buffer. When running SDS-PAGE gel, 10mM DTT is recommended.

Stability: Lyophilized Bcl-2 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Bcl-2 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Purity: Greater than 95.0% as determined by RP-HPLC and SDS-PAGE.

Applications: Input marker or positive control (Western Blotting); Functional studies.

Laboratory Reagent For Research Use Only

JK 05/02/23

Visit our website for your local distributor.

CEDARLANE®
www.cedarlanelabs.com



An ISO 9001:2015 and ISO 13485:2016
registered company.

In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020
e-mail: general@cedarlanelabs.com

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138
e-mail: service@cedarlanelabs.com