



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TECHNICALLY *Speaking*

Place your order with CEDARLANE® or your local distributor.

Please contact CEDARLANE® for lot specific information.

Mouse Anti-Human Androgen Receptor Monoclonal Antibody

CLSG36511

Lot:

Immunogen:

BALB/C mice were injected with a synthetic peptide, amino acid 299-315 (STEDTAEYSPFKGGYTK) of human androgen receptor.

Clone:

AR 441

Isotype:

IgG1

Format:

1 ml purified immunoglobulin fraction containing sodium azide as preservative.

Titer/Working Dilution:

This antibody may be diluted to a titer of 1:25-1:50 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user.

Staining Protocol:

We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user. Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 10 mM citrate buffer, pH 6.0 prior to immunostaining.

Specificity:

This antibody is specific to a protein of 110 kD, identified as androgen receptor. This antibody reacts with full length of AR and also with the newly described A form of the receptor. This antibody does not cross react with estrogen, progesterone or glucocorticoid receptors.

Positive Control:

Prostate carcinoma

Cellular Localization:

Nuclear

Other applications:

Western Blotting (1:50 to 1:100)

Storage:

Store the antibody at 4-8° c. Do not use it beyond the expiration date stated on the label.

For Research Use Only. Not for Diagnostic or Therapeutic Work.

JCr 06/15/06

For more information or to place an order please contact...

CEDARLANE®
LABORATORIES LIMITED



toll free: 1-800-268-5058
in North America

phone: (905) 878-8891 • fax: (905) 878-7800

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information
website: www.cedarlanelabs.com • e-mail: info@cedarlanelabs.com