



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TECHNICALLY *Speaking*

Place your order with CEDARLANE® or your local distributor.

Please contact CEDARLANE® for lot specific information.

Purified Mouse Anti-bcl-X (Apoptosis Marker) Monoclonal Antibody

CLSG36523

Lot:

Immunogen:

BALB/C mice were injected with a synthetic peptide of amino acid 3-14 (Cys-QSNRELVVDFLS) of human bcl-X protein.

Clone:

2H12

Isotype:

IgG2a

Format:

1 ml. This antibody is supplied as purified immunoglobulin fraction containing sodium azide as preservative.

Titer/Working Dilution:

This antibody may be diluted to a titer of 1:25-1:50 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user.

Staining Protocol:

We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user. Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 10 mM citrate buffer, pH 6.0 prior to immunostaining.

Specificity:

This antibody recognizes a 27 kD protein identified as bcl-X protein. This antibody is highly specific to bcl-X and does not show cross reactivity with bcl-2 alpha or bax protein. This antibody cross reacts with human, mouse, rat and pig.

Positive Control:

Hodgkin's lymphoma

Continued Overleaf...

For more information or to place an order please contact...

CEDARLANE®
LABORATORIES LIMITED



toll free: 1-800-268-5058
in North America

phone: (905) 878-8891 • fax: (905) 878-7800

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information
website: www.cedarlanelabs.com • e-mail: info@cedarlanelabs.com

- Cellular Localization:** Cytoplasmic, cell membrane
- Other applications:** Immunoprecipitation (1:50), Western Blotting (1:50 to 1:100)
- Storage:** Store the antibody at 4-8°C. Do not use it beyond the expiration date stated on the label.
- References:**
- i) Hsu et al. J Biol. Sci. 272: 13829, 1997.
 - ii) Hsu et al. Proc Natl Acad Sci. USA 94: 3668, 1997.
 - iii) Wolter et al. J Cell Biol. 139: 1281, 1997.

For Research Use Only. Not for Diagnostic or Therapeutic Work.