



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



TECHNICALLY *Speaking*

Place your order with CEDARLANE® or your local distributor.

Please contact CEDARLANE® for lot specific information.

Purified Rabbit anti-CD3 Zeta Polyclonal Antibody

CLSG36906
CLSG36906-05
Lot:

Immunogen:	A synthetic peptide corresponding to a.a. 36 –54 a.a. of the human CD3 zeta protein.
Host:	Rabbit
Format:	1 ml (CLSG36906) or 0.5 ml (CLSG36906-05). Purified immunoglobulin fraction of rabbit antiserum against CD3 zeta containing sodium azide as preservative.
Titer/Working Dilution:	This antibody may be diluted to a titer of 1:50-1:100 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user.
Staining Protocol:	We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation conditions should be determined by the user. <u>Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 10 mM citrate buffer, pH 6.0 prior to immunostaining.</u>
Specificity:	This antibody reacts with a 19 kDa protein. CD3 consists of five invariant polypeptide chains that associate to form three dimers: a heterodimer of gamma and epsilon chains, a heterodimer of delta and epsilon chains and a homodimer of two zeta chains or a heterodimer of zeta and eta chains. Zeta chain consists of three consecutive copies of the same motif. This antibody cross reacts with mouse and rat.
Positive Control:	Tonsil
Cellular Localization:	Cell membrane
Storage:	Store the antibody at 2-8°C. Do not use it beyond the expiration date stated on the label.

Laboratory Reagent For Research Use Only

JV 11/20/08

For more information or to place an order please contact...

CEDARLANE®
LABORATORIES LIMITED



toll free: 1-800-268-5058
in North America

phone: (905) 878-8891 • fax: (905) 878-7800

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information
website: www.cedarlanelabs.com • e-mail: info@cedarlanelabs.com