



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Technically
Speaking

CEDARLANE[®]
www.cedarlanelabs.com



Delivering You Today's Innovations for
the Science of Tomorrow™

Mouse Anti-Inhibin alpha Monoclonal Antibody

CLSG80028
CLSG8008-05
Lot: L524

- Clone:** R1
- Immuogen:** Synthetic peptide corresponding to the 1-32 amino terminal peptide of the alpha subunit of human inhibin.
- Isotype:** IgG2a
- Format:** 1 ml (**CLSG80028**) or 0.5 ml (**CLSG8008-05**). This antibody is supplied as tissue culture supernatant containing sodium azide as preservative.
- Titer/Working Dilution:** This antibody may be diluted to a titer of 1:25-1:50 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user.
- Staining Protocol:** We suggest an incubation period of 30 to 60 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user. Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 10 mM citrate buffer, pH 6.0 prior to immunostaining.
- Specificity:** Inhibin is a dimeric glycoprotein hormone from TGF-beta family made of alpha and beta subunits. This antibody recognizes the 32 kDa alpha subunit of human inhibin. Inhibin alpha is expressed in a wide variety of human tissues, such as brain, prostate, adrenal, granulosa cells of ovary, sertoli cells of testis and in fetoplacental unit. This antibody cross reacts with human and weakly with cow and pig.
- Positive Control:** Ovary
- Cellular Localization:** Cytoplasmic
- Storage:** Store the antibody at 2-8°C. Do not use it beyond the expiration date stated on the label.
- References:**
Costa et al. Human Pathol 28: 1247, 1997
Groome et al. Hybridoma 9:31, 1990
Matis-Goju et al. Human Pathol 29:840, 1998
Zheng et al. App Immunol Mol Morph 7:29, 1999

Laboratory Reagent For Research Use Only

JV 06/30/09

Visit our website for your local distributor.

CEDARLANE[®]



www.cedarlanelabs.com

An ISO 9001:2000 and ISO 13485:2003
registered company.

In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020
e-mail: general@cedarlanelabs.com

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138
e-mail: service@cedarlanelabs.com