



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TECHNICALLY *Speaking*

Place your order with CEDARLANE® or your local distributor.

Please contact CEDARLANE® for lot specific information.

Mouse Anti-p27^{Kip1} (Mitotic Inhibitor/Suppressor Protein) Monoclonal Antibody

CLSG80223-05

Lot:

Immunogen:	BALB/C mice were injected with mouse recombinant p27 protein.
Clone:	DCS-72.F6
Isotype:	IgG1
Format:	0.5 mL purified immunoglobulin fraction containing sodium azide as preservative.
Titer/Working Dilution:	This antibody may be diluted to a titer of 1:25-1:50 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user.
Staining Protocol:	We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation should be determined by the user. <u>Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 10 mM citrate buffer, pH 6.0 prior to immunostaining.</u>
Specificity:	This antibody is specific to a protein of 27 kD known as p27 ^{Kip1} a cell cycle regulatory mitotic inhibitor. This antibody recognizes an epitope between amino acids 83-204 of p27. This antibody does not show any cross reactivity with other related mitotic inhibitors.
Positive Control:	Colon carcinoma
Cellular Localization:	Nuclear
Other applications:	Immunoprecipitation (1:50), Western Blotting (1:50 to 1:100)
Storage:	Store the antibody at 2-8° c. Do not use it beyond the expiration date stated on the label.

For Research Use Only. Not for Diagnostic or Therapeutic Work.

JJ 10/11/06

For more information or to place an order please contact...

CEDARLANE®
LABORATORIES LIMITED



toll free: 1-800-268-5058
in North America

phone: (905) 878-8891 • fax: (905) 878-7800

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information
website: www.cedarlanelabs.com • e-mail: info@cedarlanelabs.com