



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TECHNICALLY *Speaking*

Place your order with CEDARLANE® or your local distributor.

Please contact CEDARLANE® for lot specific information.

Purified Rabbit anti-Caspase 3 (CPP32) Polyclonal Antibody

CLSG80436
CLSG80436-05
Lot:

Immunogen: Recombinant full-length human Caspase 3 protein.

Host: Rabbit

Format: 1 ml (CLSG80436) or 0.5 ml (CLSG80436-05), purified immunoglobulin fraction of rabbit antiserum against Caspase 3 containing sodium azide as preservative.

Titer/Working Dilution: This antibody may be diluted to a titer of 1:50-1:100 in an ABC method. However, depending on the staining conditions employed, we suggest that final dilution should be determined by the user.

Staining Protocol: We suggest an incubation period of 30 minutes at room temperature. However, depending upon the fixation conditions and the staining system employed, optimal incubation conditions should be determined by the user. Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 10 mM citrate buffer, pH 6.0 prior to immunostaining.

Specificity: This antibody reacts with active form of caspase 3 (17 kDa protein). Caspase 3 is a member of the interleukin-1 beta-converting enzyme family. Caspase 3 is thought to be associated with induction of apoptosis. Caspase 3 is synthesized as inactive 32 kDa proenzyme and is processed during apoptosis generating two subunits of 17 kDa and 12 kDa. Caspase 3 stains the epithelial cells of skin, renal proximal tubules and collecting ducts. This antibody cross-reacts with mouse, rat, monkey, rabbit, hamster, cow, sheep, pig and dog.

Positive Control: Tonsil

Cellular Localization: Cytoplasmic, some nuclear

Storage: Store the antibody at 2-8°C. Do not use it beyond the expiration date stated on the label.

Continued...

For more information or to place an order please contact...

CEDARLANE®
LABORATORIES LIMITED



toll free: 1-800-268-5058
in North America

phone: (905) 878-8891 • fax: (905) 878-7800

5516 - 8th Line, R.R.#2, Hornby, Ontario, CANADA L0P 1E0

or visit our website for a list of our international distributors including contact information
website: www.cedarlanelabs.com • e-mail: info@cedarlanelabs.com

References:

Krajewska et al. Cancer Res 57: 1605, 1997.
Mallat et al. Circulation 96: 424, 1997.

JV 03/13/09

Laboratory Reagent For Research Use Only