



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)



*Technically
Speaking*

CEDARLANE® 
www.cedarlanelabs.com

Conveniently Delivering You Today's Innovations
for the Science of Tomorrow™

Rabbit anti-CD20 Monoclonal Antibody

CLSG90504

Lot:

Intended Use:

This product is intended for qualitative immunohistochemistry with normal and neoplastic formalin-fixed, paraffin-embedded tissue sections, to be viewed by light microscopy. Clinical interpretation of staining results should be accompanied by histological studies with proper controls. Patients' clinical histories and other relevant diagnostic tests should be utilized by a qualified person(s) when evaluating and interpreting results.

Immunogen:

A synthetic peptides from the C-terminus of human CD20 protein.

Clone:

BV11

Isotype:

IgG

Size:

0.5 ml

Format:

This antibody is supplied as tissue culture supernatant and diluted in buffer pH 7.5, containing BSA and sodium azide as preservatives.

Titer/Working Dilution:

This antibody may be diluted to a titer of 1:25 to 1:50 in an ABC method. The final dilution should be determined by the user based upon the staining conditions employed.

Staining Protocol:

We suggest an incubation period of 30-60 minutes at room temperature. Optimal incubation conditions should be determined by the user based upon the fixation conditions and staining system employed.

Continued...

Visit our website for your local distributor.

CEDARLANE® 

www.cedarlanelabs.com

An ISO 9001:2000 and ISO 13485:2003
registered company.

In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020
e-mail: general@cedarlanelabs.com

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138
e-mail: service@cedarlanelabs.com

Specificity:	This antibody reacts with CD20, which is a 33kDa leukocyte surface antigen consisting of four transmembrane regions and Cytoplasmic N- and C-termini. CD20 is expressed primarily on B cells but has also been detected on both normal and neoplastic T cells. Activation of CD20 significantly increases DNA synthesis and is thought to involve basic helix-loop-helix leucine zipper transcription factors.
Positive Control:	Tonsil
Cellular Localization:	Cytoplasmic and membrane
Storage:	Store at 2-8°C. Do not use beyond the expiration date stated on the label.

Laboratory Reagent For Research Use Only

BA 05/17/12