



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Recombinant Toxoplasma Gondii P30 (SAG1)

CLTOX264
CLTOX264-2
CLTOX264-3

Introduction: The life cycle of *Toxoplasma gondii* has two phases. The coccidia like takes place only in members of the Felidae family which makes these animals the parasite's primary host. The asexual part of the life cycle can take place in any warm-blooded animal, like other mammals (including felines) and birds. *T. gondii* constructing daughter scaffolds within the mother cell. In the intermediate hosts (including felines), the parasite invades cells, forming intracellular so-called parasitophorous vacuoles containing bradyzoites, the slowly replicating form of the parasite. Vacuoles form tissue cysts mainly within the muscles and brain. Since they are within cells, the host's immune system does not detect these cysts. Resistance to antibiotics varies, but the cysts are very difficult to eradicate entirely. Within these vacuoles *T. gondii* propagates by a series of binary fissions until the infected cell eventually bursts and tachyzoites are released. Tachyzoites are the motile, asexually reproducing form of the parasite. Unlike the bradyzoites, the free tachyzoites are usually efficiently cleared by the host's immune response, although some manage to infect cells and form bradyzoites, thus maintaining the infection.

Description: The E. Coli derived recombinant protein contains the p30 (SAG1) immunodominant regions, amino acids 45-198 and fused to a 26kDa GST tag, having a total Mw of 42.3 kDa.

Source: *Escherichia Coli*.

Presentation: 100 µg (CLTOX264), 0.5 mg (CLTOX264-2), or 1 mg (CLTOX264-3), 50mM Tris-HCl pH 8, 1.5M urea and 50% glycerol.

Purification Method: *Toxoplasma* protein was purified by proprietary chromatographic technique.

Stability: *Toxoplasma* protein although stable at 4°C for 1 week, should be stored below -18°C. Please prevent freeze thaw cycles.

Purity: *Toxoplasma* protein is >95% pure as determined by 10% PAGE (Coomassie staining).

Specificity: Immunoreactive with sera of *Toxoplasma gondii*-infected individuals.

Applications: *Toxoplasma* antigen is suitable for ELISA and Western blots, excellent antigen for detection of *Toxoplasma gondii*- with minimal specificity problems.

Laboratory Reagent For Research Use Only

MW 03/04/20

Visit our website for your local distributor.

CEDARLANE®



www.cedarlanelabs.com

An ISO 9001:2015 and ISO 13485:2016
registered company.

In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020
e-mail: general@cedarlanelabs.com

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138
e-mail: service@cedarlanelabs.com