



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC Handels GmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Technically
Speaking

CEDARLANE® 
www.cedarlanelabs.com

Conveniently Delivering You Today's Innovations
for the Science of Tomorrow™

Mouse Anti-CD253/TRAIL Monoclonal Antibody

CLX106NA
CLX106PE

Clone: 2E5

Isotype: Mouse IgG1

Specificity:

The antibody 2E5 reacts with TRAIL (APO-2L), a 21 kDa cytotoxic protein, activator of rapid apoptosis in tumor cells. TRAIL is mainly expressed in spleen, lung, prostate and also in many other tissues.

Immunogen: Recombinant soluble fragment (aa 95-281) of human TRAIL.

Species Reactivity: Human.

Application:

Functional Application (CLX106NA)

The antibody 2E5 has high neutralizing activity for human TRAIL in biological assays.

Flow Cytometry (CLX106PE; CLX106NA)

Recommended dilution: 1-10 µg/ml

***Optimal working concentrations should be determined by each investigator.**

Conjugate Preparation:

PE: The purified antibody is conjugated with R-Phycoerythrin (PE) under optimum conditions. The conjugate is purified by size-exclusion chromatography.

Presentation:

Azide Free: 0.1 mg (1 mg/mL) purified IgG buffered in PBS without sodium azide, approx. pH 7.4. 0.2 µm filter sterilized. Endotoxin level is less than 0.01 EU/µg of the protein, as determined by the LAL test. (Purified from hybridoma culture supernatant by protein-A affinity chromatography).

PE: 0.1 mg (0.1 mg/mL) PE conjugated IgG buffered in PBS with 15 mM sodium azide, approx. pH 7.4.

Storage / Stability:

Store in the dark at 2-8°C. Do not freeze all formats. Avoid prolonged exposure to light of PE conjugate. Do not use after expiration date stamped on vial label.

Continued Overleaf....

Visit our website for your local distributor.

CEDARLANE® 

www.cedarlanelabs.com

An ISO 9001:2000 and ISO 13485:2003
registered company.

In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020
e-mail: general@cedarlanelabs.com

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138
e-mail: service@cedarlanelabs.com

Background:

Human CD253 / TRAIL (TNF-Related Apoptosis Inducing Ligand), also called Apo2, is a type II membrane protein from the TNF family. TRAIL is a cytotoxic protein which activates rapid apoptosis in tumor cells, but not in normal cells. TRAIL-induced apoptosis, is achieved through binding to two death-signaling receptors, DR4 (CD261 / TRAIL-R1) and DR5 (CD262 / TRAIL-R2).

References:

Plasilova M, Zivny J, Jelinek J, Neuwirtova R, Cermak J, Necas E, Andera L, Stopka T: TRAIL (Apo2L) suppresses growth of primary human leukemia and myelodysplasia progenitors. *Leukemia*. 2002 Jan;16(1):67-73.

Hyer ML, Croxton R, Krajewska M, Krajewski S, Kress CL, Lu M, Suh N, Sporn MB, Cryns VL, Zapata JM, Reed JC: Synthetic triterpenoids cooperate with tumor necrosis factor-related apoptosis-inducing ligand to induce apoptosis of breast cancer cells. *Cancer Res*. 2005 Jun 1;65(11):4799-808.

Laboratory Reagent For Research Use Only

JK 07/05/17