



**SZABO  
SCANDIC**

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](http://linkedin.com/company/szaboscandic)



## Mouse Anti-CD8 Monoclonal Antibody

CLX10AP	CLX10PE
CLX10B	CLX10PCP
CLX10F	CLX10APC

**Clone:** MEM-31

**Isotype:** Mouse IgG2a

**Specificity:**

The antibody MEM-31 recognizes a conformationally-dependent epitope of CD8, a cell surface glycoprotein found on most cytotoxic T lymphocytes that mediates efficient cell-cell interactions within the immune system. CD8 is a disulfide-linked dimer and exists as a CD8 alpha/alpha homodimer or CD8 alpha/beta heterodimer (each monomer approx. 32-34 kDa). The antibody does not react with formaldehyde-fixed cells; negative in Western Blotting application. HLDA III; WS Code T 575

**Immunogen:** Crude thymus membrane fraction.

**Species Reactivity:** Human.

**Application:** Flow Cytometry, Immunoprecipitation and Mass Cytometry.

**Conjugate Preparation:**

The purified antibody is conjugated with Biotin-LC-NHS, Fluorescein isothiocyanate (FITC), R-Phycoerythrin (PE), Peridinin-chlorophyll-protein complex (PerCP) or cross-linked Allophycocyanin (APC) under optimum conditions. The conjugates are purified by size-exclusion chromatography and adjusted for direct use (FITC, PE, APC, PerCP). No reconstitution is necessary.

**Presentation:**

**Purified:** 0.1 mg (1 mg/mL) purified IgG buffered in PBS with 15 mM sodium azide, approx. pH 7.4. (Purified by protein-A affinity chromatography; purity > 95% by SDS-PAGE).

**Biotin:** 0.1 mg (1 mg/mL) of Biotin conjugated IgG buffered in tris buffered saline (TBS) with 15 mM sodium azide, approx. pH 8.0.

**FITC:** 2 mL of FITC conjugated IgG buffered in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide. Sufficient for 100 tests.

**PE:** 2 mL of PE conjugated IgG buffered in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide. Sufficient for 100 tests.

**PerCP:** 1 mL of APC conjugated IgG buffered in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide. Sufficient for 100 tests.

**APC:** 1 mL of APC conjugated IgG buffered in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide. Sufficient for 100 tests.

*Continued Overleaf.....*

Visit our website for your local distributor.

**CEDARLANE®**  
  
[www.cedarlanelabs.com](http://www.cedarlanelabs.com)

An ISO 9001:2000 and ISO 13485:2003  
registered company.

**In CANADA: Toll Free: 1-800-268-5058**

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020  
e-mail: general@cedarlanelabs.com

**In the USA: Toll Free: 1-800-721-1644**

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138  
e-mail: service@cedarlanelabs.com

**Storage / Stability:**

Store in the dark at 2-8°C. Do not freeze all formats. Avoid prolonged exposure to light of conjugates. Do not use after expiration date stamped on vial label.

**Usage:**

Recommended dilutions for Flow Cytometry analysis of human blood cells:

**Purified:** 1 µg/ml

**Biotin:** 1:2500 dilution

**FITC:** 20 µl reagent / 100 µl of whole blood or  $10^6$  cells in a suspension.

**PE:** 20 µl reagent / 100 µl of whole blood or  $10^6$  cells in a suspension.

**PerCP:** 10 µl reagent / 100 µl of whole blood or  $10^6$  cells in a suspension.

**APC:** 10 µl reagent / 100 µl of whole blood or  $10^6$  cells in a suspension.

**\*Optimal working concentrations should be determined by the investigator.**

**Background:**

The CD8 T cell coreceptor (monomer approx. 32-34 kDa) is expressed as alpha/beta heterodimer on majority of MHC I-restricted conventional T cells and thymocytes and as alpha/alpha homodimer on subsets of memory T cells, intraepithelial lymphocytes, NK cells and dendritic cells. Regulation of CD8 beta level on T cell surface seems to be an important mechanism to control their effector function. Assembly of CD8 alpha-beta but not alpha-alpha dimers is connected with formation or localization to the lipid rafts. Recruiting triggered TCR complexes to these membrane microdomains as well as affinity of TCR to MHC I is modulated by CD8, thereby affecting the functional diversity of the TCR signaling.

**References:**

- \*Devine L, Thakral D, Nag S, Dobbins J, Hodsdon ME, Kavathas PB: Mapping the binding site on CD8 beta for MHC class I reveals mutants with enhanced binding. *J Immunol.* 2006 Sep 15;177(6):3930-8.
- \*Pang DJ, Hayday AC, Bijlmakers MJ.: CD8 Raft localization is induced by its assembly into CD8alpha beta heterodimers, Not CD8alpha alpha homodimers. *J Biol Chem.* 2007 May 4;282(18):13884-94.
- \*van den Berg HA, Wooldridge L, Laugel B, Sewell AK: Coreceptor CD8-driven modulation of T cell antigen receptor specificity. *J Theor Biol.* 2007 Nov 21;249(2):395-408.
- \*Horejsí V et al.: Monoclonal antibodies against human leucocyte antigens. I. Antibodies against beta-2-microglobulin, immunoglobulin kappa light chains, HLA-DR-like antigens, T8 antigen, T1 antigen, a monocyte antigen, and a pan-leucocyte antigen. *Folia Biol. (Praha)* 32, 12 (1986).
- \*Leukocyte Typing III., McMichael A. J. et al (Eds.), Oxford University Press (1987).
- \*Horejsí V, Angelisová P, Bazil V, Kristofová H, Stoyanov S, Stefanová I, Hausner P, Vosecký M, Hilgert I: Monoclonal antibodies against human leucocyte antigens. II. Antibodies against CD45 (T200), CD3 (T3), CD43, CD10 (CALLA), transferrin receptor (T9), a novel broadly expressed 18-kDa antigen (MEM-43) and a novel antigen of restricted expression (MEM-74). *Folia Biol (Praha)*. 1988;34(1):23-34.
- \*Brdicková N, Brdicka T, Angelisová P, Horváth O, Spicka J, Hilgert I, Paces J, Simeoni L, Kliche S, Merten C, Schraven B, Horejsí V: LIME: a new membrane Raft-associated adaptor protein involved in CD4 and CD8 coreceptor signaling. *J Exp Med.* 2003 Nov 17;198(10):1453-62.
- \*Drbal K, Moertelmaier M, Holzhauser C, Muhammad A, Fuertbauer E, Howorka S, Hinterberger M, Stockinger H, Schütz GJ: Single-molecule microscopy reveals heterogeneous dynamics of lipid raft components upon TCR engagement. *Int Immunol.* 2007 May;19(5):675-84.
- \*Estefanía E, Flores R, Gómez-Lozano N, Aguilar H, López-Botet M, Vilches C: Human KIR2DL5 is an inhibitory receptor expressed on the surface of NK and T lymphocyte subsets. *J Immunol.* 2007 Apr 1;178(7):4402-10.
- \*Linnebacher M, Wienck A, Boeck I, Klar E: Identification of an MSI-H tumor-specific cytotoxic T cell epitope generated by the (-1) frame of U79260(FTO). *J Biomed Biotechnol.* 2010;2010:841451.
- \*Kanderova V, Kuzilkova D, Stuchly J, Vaskova M, Brdicka T, Fiser K, Hrusak O, Lund-Johansen F, Kalina T: High-resolution Antibody Array Analysis of Childhood Acute Leukemia Cells. *Mol Cell Proteomics.* 2016 Apr;15(4):1246-61.

**Laboratory Reagent For Research Use Only**