



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



FITC Mouse Anti-CD36 Monoclonal Antibody

CLX461F

Lot:

Size: 100 tests

Clone: CB38 (NL07)

Isotype: Mouse IgM

Regulatory Status: RUO

Specificity: The mouse monoclonal antibody CB38 (NL07) recognizes CD36 (GPIIIb), a 85-113 kDa integral membrane glycoprotein expressed on platelets, macrophages, endothelial cells, early erythroid cells and megakaryocytes.

HLDA IV.; WS Code P106

Immunogen: Living human myeloid cells

Species Reactivity: Human

Preparation: The purified antibody is conjugated with Fluorescein isothiocyanate (FITC) under optimum conditions. The reagent is free of unconjugated FITC and adjusted for direct use. No reconstitution is necessary.

Storage Buffer: The reagent is provided in phosphate buffered saline (PBS) containing 15 mM sodium azide and 0.2% (w/v) high-grade protease free Bovine Serum Albumin (BSA) as a stabilizing agent.

Storage/Stability: Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label. Short-term exposure to room temperature should not affect the quality of the reagent. However, if reagent is stored under any conditions other than those specified, the conditions must be verified by the user.

Usage: The reagent is designed for Flow Cytometry analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or 10⁶ cells in a suspension. The content of a vial (0.4 ml) is sufficient for 100 tests.

Continued...

Visit our website for your local distributor.

CEDARLANE®

www.cedarlanelabs.com

An ISO 9001:2000 and ISO 13485:2003
registered company.

In CANADA: Toll Free: 1-800-268-5058

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020
e-mail: general@cedarlanelabs.com

In the USA: Toll Free: 1-800-721-1644

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138
e-mail: service@cedarlanelabs.com

Background:

CD36 (fatty acid translocase, FAT) is an 85-113 kDa ditopic glycosylated protein that belongs to the class B family of scavenger receptors. CD36 is expressed by most resting marginal zone B cells but not by follicular and B1 B cells, and it is rapidly induced on Follicular B cells in vitro upon TLR and CD40 stimulation. CD36 does not affect the development of B cells, but modulates both primary and secondary antibody response. Similarly to glucose transporter GLUT4, CD36 is translocated from intracellular pools to the plasma membrane following cell stimulation by insulin. In mouse, CD36 is responsible for gustatory perception of long-chain fatty acids.

References:

- *Leukocyte Typing IV., Knapp W. et al. (Eds.), Oxford University Press (1989).
- *Alessio M, Ghigo D, Garbarino G, Geuna M, Malavasi F: Analysis of the human CD36 leucocyte differentiation antigen by means of the monoclonal antibody NL07. *Cell Immunol.* 1991 Oct 15;137(2):487-500.
- *Alessio M, Roggero S, Bussolino F, Saitta M, Malavasi F: Characterization of the murine monoclonal antibody NL07 specific for the human thrombospondin receptor (CD36 molecule). *Curr Stud Hematol Blood Transfus.* 1991;(58):182-6.
- *Alessio M, Greco NJ, Primo L, Ghigo D, Bosia A, Tandon NN, Ockenhouse CF, Jamieson GA, Malavasi F: Platelet activation and inhibition of malarial cytoadherence by the anti-CD36 IgM monoclonal antibody NL07. *Blood.* 1993 Dec 15;82(12):3637-47.
- *Fadok VA, Warner ML, Bratton DL, Henson PM: CD36 is required for phagocytosis of apoptotic cells by human macrophages that use either a phosphatidylserine receptor or the vitronectin receptor (alpha v beta 3). *J Immunol.* 1998 Dec 1;161(11):6250-7.
- *Matasić R, Dietz AB, Vuk-Pavlović S: Dexamethasone inhibits dendritic cell maturation by redirecting differentiation of a subset of cells. *J Leukoc Biol.* 1999 Dec;66(6):909-14.
- *Hoffmann PR, deCathelineau AM, Ogden CA, Leverrier Y, Bratton DL, Daleke DL, Ridley AJ, Fadok VA, Henson PM: Phosphatidylserine (PS) induces PS receptor-mediated macropinocytosis and promotes clearance of apoptotic cells. *J Cell Biol.* 2001 Nov 12;155(4):649-59.
- *Kapinsky M, Torzewski M, Büchler C, Duong CQ, Rothe G, Schmitz G: Enzymatically degraded LDL preferentially binds to CD14(high) CD16(+) monocytes and induces foam cell formation mediated only in part by the class B scavenger-receptor CD36. *Arterioscler Thromb Vasc Biol.* 2001 Jun;21(6):1004-10.
- *McKenna RW, Washington LT, Aquino DB, Picker LJ, Kroft SH: Immunophenotypic analysis of hematogones (B-lymphocyte precursors) in 662 consecutive bone marrow specimens by 4-color flow cytometry. *Blood.* 2001 Oct 15;98(8):2498-507.
- *Moniuszko M, Kowal K, Rusak M, Pietruck M, Dabrowska M, Bodzenta-Lukaszyk A: Monocyte CD163 and CD36 expression in human whole blood and isolated mononuclear cell samples: influence of different anticoagulants. *Clin Vaccine Immunol.* 2006 Jun;13(6):704-7.
- *And many other.

Laboratory Reagent For Research Use Only

BA 11/19/13