



**SZABO  
SCANDIC**

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](http://linkedin.com/company/szaboscandic)



## APC Mouse Anti-Kappa Light Chains Monoclonal Antibody

**CLX465APC**

**Lot:**

**Size:** 100 tests

**Clone:** TB28-2

**Isotype:** Mouse IgG1

**Regulatory Status:** RUO

**Specificity:** The antibody TB28-2 recognizes human Ig kappa light chains of both secreted and cell surface immunoglobulin. It detects also free kappa light chains.

**Immunogen:** Human IgG-kappa myeloma protein

**Species Reactivity:** Human

**Preparation:** The purified antibody is conjugated with cross-linked Allophycocyanin (APC) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.

**Storage Buffer:** The reagent is provided in phosphate buffered saline (PBS) containing 15 mM sodium azide and 0.2% (w/v) high-grade protease free Bovine Serum Albumin (BSA) as a stabilizing agent.

**Storage/Stability:** Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label. Short-term exposure to room temperature should not affect the quality of the reagent. However, if reagent is stored under any conditions other than those specified, the conditions must be verified by the user.

**Usage:** The reagent is designed for Flow Cytometry analysis of human blood cells using 10 µl reagent / 100 µl of whole blood or 10<sup>6</sup> cells in a suspension. The content of a vial (1 ml) is sufficient for 100 tests.

*Continued...*

Visit our website for your local distributor.

**CEDARLANE®**  
  
[www.cedarlanelabs.com](http://www.cedarlanelabs.com)

An ISO 9001:2000 and ISO 13485:2003  
registered company.

**In CANADA: Toll Free: 1-800-268-5058**

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020  
e-mail: general@cedarlanelabs.com

**In the USA: Toll Free: 1-800-721-1644**

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138  
e-mail: service@cedarlanelabs.com

**Background:**

Immunoglobulin classes share the same basic four polypeptide chain structure of two heavy chains (five heavy chains types) and two light chains (kappa, lambda; both having a molecular weight of 22.5kDa). Kappa and lambda consist of a variable region and a constant region and can easily be differentiated by the antigenic properties of the constant region. The ratio of kappa to lambda is 70:30.

**References:**

- \*Kiyotaki M, Cooper MD, Bertoli LF, Kearney JF, Kubagawa H: Monoclonal anti-Id antibodies react with varying proportions of human B lineage cells. *J Immunol*. 1987 Jun 15;138(12):4150-8.
- \*Nakamura T, Kubagawa H, Cooper MD: Heterogeneity of immunoglobulin-associated molecules on human B cells identified by monoclonal antibodies. *Proc Natl Acad Sci U S A*. 1992 Sep 15;89(18):8522-6.
- \*Böttcher S, Ritgen M, Buske S, Gesk S, Klapper W, Hoster E, Hiddemann W, Unterhalt M, Dreyling M, Siebert R, Kneba M, Pott C; EU MCL MRD Group: Minimal residual disease detection in mantle cell lymphoma: methods and significance of four-color flow cytometry compared to consensus IGH-polymerase chain reaction at initial staging and for follow-up examinations. *Haematologica*. 2008 Apr;93(4):551-9.
- \*Karandikar NJ, Aquino DB, McKenna RW, Kroft SH: Transient myeloproliferative disorder and acute myeloid leukemia in Down syndrome. An immunophenotypic analysis. *Am J Clin Pathol*. 2001 Aug;116(2):204-10.
- \*Jourdan M, Caraux A, De Vos J, Fiol G, Larroque M, Cognot C, Bret C, Duperray C, Hose D, Klein B: An in vitro model of differentiation of memory B cells into plasmablasts and plasma cells including detailed phenotypic and molecular characterization. *Blood*. 2009 Dec 10;114(25):5173-81.

**Laboratory Reagent For Research Use Only**

BA 01/22/14