



**SZABO  
SCANDIC**

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](http://linkedin.com/company/szaboscandic)





Conveniently Delivering You Today's Innovations  
for the Science of Tomorrow™

## FITC Mouse Anti-Human CD63 Monoclonal Antibody

### CLX66F

**Size:** 100 Tests

**Clone:** MEM-259

**Isotype:** Mouse IgG1

**Specificity:** The antibody MEM-259 reacts with CD63 (LAMP-3), a 40-60 kDa tetraspan glycoprotein expressed by granulocytes, platelets, T cells, monocytes/macrophages and endothelial cells. Cell surface exposition of CD63 is usually activation-dependent.

**Immunogen:** HPB-ALL T cell line

**Species Reactivity:** Human

**Preparation:** The purified antibody is conjugated with Fluorescein isothiocyanate (FITC) under optimum conditions. The reagent is free of unconjugated FITC and adjusted for direct use. No reconstitution is necessary.

**Storage Buffer:** The reagent is provided in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide.

**Storage / Stability:** Store in the dark at 2-8°C. Do not freeze. Avoid prolonged exposure to light. Do not use after expiration date stamped on vial label.

**Usage:** The reagent is designed for Flow Cytometry analysis of human blood cells using 20 µl reagent / 100 µl of whole blood or 10<sup>6</sup> cells in a suspension. The content of a vial (2 ml) is sufficient for 100 tests.

*Continued Overleaf.....*

Visit our website for your local distributor.

**CEDARLANE®**

[www.cedarlanelabs.com](http://www.cedarlanelabs.com)

An ISO 9001:2000 and ISO 13485:2003  
registered company.

**In CANADA: Toll Free: 1-800-268-5058**

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020  
e-mail: general@cedarlanelabs.com

**In the USA: Toll Free: 1-800-721-1644**

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138  
e-mail: service@cedarlanelabs.com

**Background:**

**CD63** (LAMP-3, lysosome-associated membrane protein-3), a glycoprotein of tetraspanin family, is present in late endosomes, lysosomes and secretory vesicles of various cell types. It is also present in the plasma membrane, usually following cell activation. Hence, it has become a widely used basophil activation marker. In mast cells, however, CD63 exposition does not need their activation. CD63 interacts with integrins and affects phagocytosis and cell migration, it is also involved in H/K-ATPase trafficking regulation of ROMK1 channels. CD63 also serves as a T-cell co-stimulation molecule. Expression of CD63 can be used for predicting the prognosis in earlier stages of carcinomas.

**References:**

- \*Grützkau A, Smorodchenko A, Lippert U, Kirchhof L, Artuc M, Henz BM: LAMP-1 and LAMP-2, but not LAMP-3, are reliable markers for activation-induced secretion of human mast cells. *Cytometry A*. 2004 Sep;61(1):62-8.
- \*Mantegazza AR, Barrio MM, Moutel S, Bover L, Weck M, Brossart P, Teillaud JL, Mordoh J: CD63 tetraspanin slows down cell migration and translocates to the endosomal-lysosomal-MIICs route after extracellular stimuli in human immature dendritic cells. *Blood*. 2004 Aug 15;104(4):1183-90.
- \*Pfistershamer K, Majdic O, Stöckl J, Zlabinger G, Kirchberger S, Steinberger P, Knapp W: CD63 as an activation-linked T cell costimulatory element. *J Immunol*. 2004 Nov 15;173(10):6000-8.
- \*Israels SJ, McMillan-Ward EM: CD63 modulates spreading and tyrosine phosphorylation of platelets on immobilized fibrinogen. *Thromb Haemost*. 2005 Feb;93(2):311-8.
- \*Kwon MS, Shin SH, Yim SH, Lee KY, Kang HM, Kim TM, Chung YJ: CD63 as a biomarker for predicting the clinical outcomes in adenocarcinoma of lung. *Lung Cancer*. 2007 Jul;57(1):46-53.
- \*Lin D, Kamsteeg EJ, Zhang Y, Jin Y, Sterling H, Yue P, Roos M, Duffield A, Spencer J, Caplan M, Wang WH: Expression of tetraspan protein CD63 activates protein tyrosine kinase (PTK) and enhances the PTK-induced inhibition of ROMK channels. *J Biol Chem*. 2008 Jan 22
- \*Cerny J, Feng Y, Yu A, Miyake K, Borgonovo B, Klumperman J, Meldolesi J, McNeil PL, Kirchhausen T: The small chemical vacuolin-1 inhibits Ca<sup>(2+)</sup>-dependent lysosomal exocytosis but not cell resealing. *EMBO Rep*. 2004 Sep;5(9):883-8. Erratum in: *EMBO Rep*. 2005 Sep;6(9):898.
- \*Heneberg P, Riegerová K, Kučera P: Pimecrolimus Is a Potent Inhibitor of Allergic Reactions to Hymenopteran Venom Extracts and Birch Pollen Allergen In Vitro. *PLoS One*. 2015 Nov 12;10(11):e0142953.

**Laboratory Reagent For Research Use Only**

JK 07/17/18