



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

GFH109 Recombinant Human TGF-β3

Description

Transforming Growth Factors (TGFs) are multifunctional peptides that regulate growth and differentiation in most cell types. The TGF-β family of proteins signal through serine/threonine kinase receptors. TGF-β isoforms (TGF-β1, -β 2, and -β 3) have overlapping, yet distinct biological actions in developing and adult tissues. TGF-β3 is an important factor in regulating cell adhesion and accelerating wound repair. TGF-β3 also functions during osteoblast proliferation, chemotaxis, and collagen synthesis.

Length	113 / 226 aa
Molecular Weight	12.9 / 25.7 kDa
Source	E. coli
Accession Number	P10600
Purity	≥95% determined by reducing and non-reducing SDS-PAGE


Specifications

Alternative Names	Transforming Growth Factor β 3, TGF β 3, TGF-β-3, TGFB3, TGFβ3
Biological Activity	Human TGF-β3 is fully biologically active when compared to standard. The activity is determined by the inhibition of IL-4 induced HT-2 cell proliferation and it is typically less than 1 ng/ml. This corresponds to an expected specific activity of 1.0 x 10 ⁶ units/mg.
Endotoxin Level	≤1.00 EU/μg as measured by kinetic LAL
Formulation	In solution on 10 mM acetic acid and 20% ethanol at a concentration of 0.25 mg/ml
AA Sequence	MALDNTNYCFR NLEENCCVRP LYIDFRQDLG WKWVHEPKGY YANFCSGPCP YLRSADTTHS TVLGLYNTLN PEASASPCCV PQDLEPLTIL YYVGRTPKVE QLSNMVVKSC KCS


Preparation and Storage

Reconstitution	Not applicable
Stability and Storage	12 months from date of receipt when stored at 4°C as supplied.

Data



Inhibition of IL-4 induced HT-2 cell proliferation for Human TGF-β3. Cell proliferation was measured to calculate the ED50, which is as expected less than 1 ng/ml.



Non-reducing (-) and reducing (+) conditions in a 4 - 20% Tris-Glycine gel stained with Coomassie Blue. 1 μg of protein was loaded in each lane. Human TGF-β3 has a predicted Mw of 25.7 kDa (each monomer is 12.9 kDa).