



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

GFH164 Recombinant Human Procalcitonin

Description

Procalcitonin is a precursor of the peptide hormone calcitonin. Procalcitonin is mainly produced by the neuroendocrine cells of the lung and thyroid gland, but may be secreted ubiquitously during inflammation or infection. Under normal expression conditions, procalcitonin is immediately cleaved into three specific fragments: calcitonin, katalcalin, and an N terminal residue. Levels of unprocessed procalcitonin are significantly increased after bacterial infection, inflammation, trauma, or shock.

Length	116 aa
Molecular Weight	12.8 kDa
Source	E. coli
Accession Number	P01258
Purity	≥95% determined by reducing and non-reducing SDS-PAGE


Specifications

Alternative Names	PCT
Biological Activity	Human Procalcitonin is fully biologically active when compared to standard. The activity can be tested using Biomerieux reagents on the Min-Vidas at 80 ng/ml with a typical recovery of 80%. There is no data currently available.
Endotoxin Level	≤1.00 EU/μg as measured by kinetic LAL
Formulation	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA)
AA Sequence	APFRSALESS PADPATLSED EARLLLAALV QDYVQMKASE LEQEQEREGS SLDSPRSKRC GNLSTCMLGT YTQDFNKFHT FPQTAIGVGA PGKKRDMSSD LERDHRPHVS MPQANAN

Preparation and Storage

Reconstitution	Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at 0.1 mg/ml, which can be further diluted into other aqueous solutions.
Stability and Storage	12 months from date of receipt when stored at -20°C to -80°C as supplied. 1 month when stored at 4°C after reconstituting as directed. 3 months when stored at -20°C to -80°C after reconstituting as directed.

Data



Non-reducing (-) and reducing (+) conditions in a 4 - 20% Tris-Glycine gel stained with Coomassie Blue. 1 μg of protein was loaded in each lane. Human Procalcitonin has a predicted Mw of 12.8 kDa.