



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## GFM52 Recombinant Mouse VEGF-120

### Description

Vascular Endothelial Growth Factor 120 (VEGF-120) is produced by a wide variety of cell types, including tumor and vascular cells. VEGF-120 is a mediator of vascular growth, vascular permeability, and plays a role in stimulating vasodilation via nitric oxide-dependent pathways. Mouse VEGF-A has several alternatively spliced isoforms, with one being VEGF-120. VEGF-120 is an angiogenic factor that is expressed throughout endochondral bone development and is important during skeletogenesis.

<b>Length</b>	121 / 242 aa
<b>Molecular Weight</b>	14.1 / 28.3 kDa
<b>Source</b>	E. coli
<b>Accession Number</b>	P15692-9
<b>Purity</b>	≥95% determined by reducing and non-reducing SDS-PAGE


### Specifications

<b>Alternative Names</b>	Vascular Endothelial Growth Factor, VEGF121, VEGF
<b>Biological Activity</b>	Mouse VEGF-120 is fully biologically active when compared to standard. The activity is determined by the proliferation of HUVEC cells and it is typically less than 5 ng/ml. This corresponds to an expected specific activity of $2.0 \times 10^5$ units/mg.
<b>Endotoxin Level</b>	≤1.00 EU/μg as measured by kinetic LAL
<b>Formulation</b>	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA)
<b>AA Sequence</b>	MPMAEGGGQN HHEVVKFMDV YQRSYCHPIE TLVDIFQEYP DEIEYIFKPS CVPLMRCGGC CNDEGLECVPT TEESNITMQI MRIKPHQGQH IGEMSFLLQHN KCECRPKKDR ARQENCDKPR R

### Preparation and Storage

<b>Reconstitution</b>	Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at 0.1 mg/ml, which can be further diluted into other aqueous solutions.
<b>Stability and Storage</b>	12 months from date of receipt when stored at -20°C to -80°C as supplied. 1 month when stored at 4°C after reconstituting as directed. 3 months when stored at -20°C to -80°C after reconstituting as directed.

### Data



Induced proliferation of HUVEC cells for Mouse VEGF-120. Cell proliferation was measured to calculate the ED50.