



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com


[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 




Human IgG Fc Monoclonal Antibody

Product Code	CSB-MA928050A0m
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Immunogen	Recombinant Human FC protein
Raised In	Mouse
Species Reactivity	Homo sapiens (Human)
Tested Applications	ELISA
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Purification Method	>95%, Protein G purified
Isotype	IgG1
Clonality	Monoclonal
Product Type	Monoclonal Antibody
Research Area	Others
Target Names	FC
Clone No.	5C7G3


Image



Human IgG Fc antibody on SDS-PAGE under reducing (R) condition. The purity of the protein is greater than 90%.




The Binding Activity of hlgG2 with Anti-Human IgG Fc antibody
Activity: Measured by its binding ability in a functional ELISA. Immobilized hlgG2 at 2 µg/mL can bind Anti-FC antibody, the EC₅₀ is 7.384 to 9.177 ng/mL.



The Binding Activity of hlgG4 with Anti-Human IgG Fc antibody

Activity: Measured by its binding ability in a functional ELISA. Immobilized hlgG4 at 2 µg/mL can bind Anti-FC antibody, the EC₅₀ is 13.30 to 19.59 ng/mL.



The Binding Activity of hlgG1 with Anti-Human IgG Fc antibody

Activity: Measured by its binding ability in a functional ELISA. Immobilized hlgG1 at 2 µg/mL can bind Anti-FC antibody, the EC₅₀ is 7.604 to 9.415 ng/mL.

Usage

For Research Use Only. Not for use in diagnostic or therapeutic procedures.